

A MODEL OF SECOND-ORDER ARITHMETIC SATISFYING AC BUT NOT DC

SY-DAVID FRIEDMAN AND VICTORIA GITMAN

ABSTRACT. We show that there is a β -model of second-order arithmetic in which the choice scheme holds, but the dependent choice scheme fails for a Π_2^1 -assertion, confirming a conjecture of Steve Simpson. We obtain as a corollary that the Reflection Principle, stating that every formula reflects to a transitive set, can fail in models of ZFC^- .

1. INTRODUCTION

Models of arithmetic are two-sorted structures, having two types of objects, which we think of as numbers and sets of numbers. Their properties are formalized using a two-sorted logic with separate variables and quantifiers for numbers and sets. By convention, we will denote number variables by lower-case letters and sets variables by upper-case letters. The language of second-order arithmetic is the language of first-order arithmetic $\mathcal{L}_A = \{+, \cdot, <, 0, 1\}$ together with a membership relation \in between numbers and sets. A multitude of second-order arithmetic theories, as well as the relationships between them, have been extensively studied (see [Sim09]).

An example of a weak second-order arithmetic theory is ACA_0 , whose axioms consist of the modified Peano axioms, where instead of the induction scheme we have the single second-order induction axiom

$$\forall X[(0 \in X \wedge \forall n(n \in X \rightarrow n + 1 \in X)) \rightarrow \forall n(n \in X)],$$

and the *comprehension scheme* for first-order formulas. The latter is a scheme of assertions stating for every first-order formula, possibly with class parameters, that there is a set whose elements are exactly the numbers satisfying the formula. One of the strongest second-order arithmetic theories is Z_2 , often referred to as *full second-order arithmetic*, which strengthens comprehension for first-order formulas in ACA_0 to full comprehension for all second-order assertions. This means that for a formula with any number of second-order quantifiers, there is a set whose elements are exactly the numbers satisfying the formula. The reals of any model of ZF is a model of Z_2 . We can further strengthen the theory Z_2 by adding choice principles for sets: the choice scheme and the dependent choice scheme.

The *choice scheme* is a scheme of assertions, which states for every second-order formula $\varphi(n, X, A)$ with a set parameter A that if for every number n , there is a set X witnessing $\varphi(n, X, A)$, then there is a single set Y collecting witnesses for every n , in the sense that $\varphi(n, Y_n, A)$ holds, where $Y_n = \{m \mid \langle n, m \rangle \in Y\}$ and $\langle n, m \rangle$ is

The first author wishes to acknowledge the support of the Austrian Science Fund (FWF) through Research Project I 1238.

any standard coding of pairs. More precisely, an instance of the choice scheme for the formula $\varphi(n, X, A)$ is

$$\forall n \exists X \varphi(n, X, A) \rightarrow \exists Y \forall n \varphi(n, Y_n, A).$$

We will denote by Σ_n^1 -AC the fragment of the choice scheme for Σ_n^1 -assertions, making an analogous definition for Π_n^1 , and we will denote the full choice scheme by Σ_∞^1 -AC. The reals of any model of $\text{ZF} + \text{AC}_\omega$ (countable choice) satisfy $\text{Z}_2 + \Sigma_\infty^1$ -AC. It is a folklore result, going back possibly to Mostowski, that the theory $\text{Z}_2 + \Sigma_\infty^1$ -AC is bi-interpretable with the theory ZFC^- (ZFC without the powerset axiom, with Collection instead of Replacement) together with the statement that every set is countable.

The *dependent choice scheme* is a scheme of assertions, which states for every second-order formula $\varphi(X, Y, A)$ with set parameter A that if for every set X , there is a set Y witnessing $\varphi(X, Y, A)$, then there is a single set Z making infinitely many dependent choices according to φ . More precisely, an instance of the dependent choice scheme for the formula $\varphi(X, Y, A)$ is

$$\forall X \exists Y \varphi(X, Y, A) \rightarrow \exists Z \forall n \varphi(Z_n, Z_{n+1}, A).$$

We will denote by Σ_n^1 -DC the dependent choice scheme for Σ_n^1 -assertions, with an analogous definition for Π_n^1 , and we will denote the full dependent choice scheme by Σ_∞^1 -DC. The reals of a model of $\text{ZF} + \text{DC}$ (dependent choice) satisfy $\text{Z}_2 + \Sigma_\infty^1$ -DC.

It is not difficult to see that the theory Z_2 implies Σ_2^1 -AC, the choice scheme for Σ_2^1 -assertions. Models of Z_2 can build their own version of Gödel's constructible universe L . If a model of Z_2 believes that a set Γ is a well-order, then it has a set coding a set-theoretic structure constructed like L along the well-order Γ . It turns out that models of Z_2 satisfy a version of Shoenfield's absoluteness with respect to their constructible universes. For every Σ_2^1 -assertion φ , a model of Z_2 satisfies φ if and only if its constructible universe satisfies φ with set quantifiers naturally interpreted as ranging over the reals. All of the above generalizes to constructible universes $L[A]$ relativized to a set parameter A .¹ Thus, given a Σ_2^1 -assertion $\varphi(n, X, A)$ for which the model satisfies $\forall n \exists X \varphi(n, X, A)$, the model can go to its constructible universe $L[A]$ to pick the least witness X for $\varphi(n, X, A)$ for every n , because $L[A]$ agrees when φ is satisfied, and then put the witnesses together into a single set using comprehension. So long as the unique witnessing set can be obtained for each n , comprehension suffices to obtain a single set of witnesses. How much more of the choice scheme follows from Z_2 ? The reals of the classical Feferman-Lévy model of ZF (see [Lév70], Theorem 8), in which \aleph_1 is a countable union of countable sets, is a β -model of Z_2 in which Σ_2^1 -AC fails. This is a particularly strong failure of the choice scheme because, as we explain below, β -models are meant to strongly resemble the full standard model $P(\omega)$.

There are two ways in which a model of second-order arithmetic can resemble the full standard model $P(\omega)$. A model of second-order arithmetic is called an ω -*model* if its first-order part is ω , and it follows that its second-order part is some subset of $P(\omega)$. But even an ω -model can poorly resemble $P(\omega)$ because it may be wrong about well-foundedness by missing ω -sequences. An ω -model of second-order arithmetic which is correct about well-foundedness is called a β -*model*. The reals of any transitive ZF -model is a β -model of Z_2 . One advantage to having a β -model of

¹Indeed, the much weaker theory ATR_0 suffices for everything we have said so far. See [Sim09] (Section VII.3 and VII.4) for details.

Z_2 is that the constructible universe it builds internally is isomorphic to an initial segment L_α of the actual constructible universe L .

The theory Z_2 also implies Σ_2^1 -DC (see [Sim09], Theorem VII.9.2), the dependent choice scheme for Σ_2^1 -assertions. In this article, we construct a symmetric submodel of a forcing extension of L whose reals form a model of second-order arithmetic in which Z_2 together with Σ_∞^1 -AC holds, but Π_2^1 -DC fails. The forcing notion we use is a tree iteration of a forcing for adding a real due to Jensen.

Jensen’s forcing, which we will call here \mathbb{P}^J , introduced by Jensen in [Jen70], is a subposet of Sacks forcing constructed in L using the \diamond principle. The poset \mathbb{P}^J has the ccc and adds a unique generic real over L . The collection of all L -generic reals for \mathbb{P}^J in any model is Π_2^1 -definable. Jensen used his forcing to show that it is consistent with ZFC that there is a Σ_3^1 -definable non-constructible real [Jen70]. Recently Lyubetsky and Kanovei extended the “uniqueness of generic filters” property of Jensen’s forcing to finite-support products of \mathbb{P}^J [KL]. They showed that in a forcing extension $L[G]$ by the ω -length finite support-product of \mathbb{P}^J , the only L -generic reals for \mathbb{P}^J are the slices of the generic filter G . The result easily extends to ω_1 -length finite support-products as well.

We in turn extend the “uniqueness of generic filters” property to tree iterations of Jensen’s forcing. We first define finite iterations \mathbb{P}_n^J of Jensen’s forcing \mathbb{P}^J , and then define an iteration of \mathbb{P}^J along a tree \mathcal{T} to be a forcing whose conditions are functions from a finite subtree of \mathcal{T} into $\bigcup_{n < \omega} \mathbb{P}_n^J$ such that nodes on level n get mapped to elements of the n -length iteration \mathbb{P}_n^J and conditions on higher nodes extend conditions on lower nodes. The functions are ordered by extension of domain and strengthening on each coordinate. We show that in a forcing extension $L[G]$ by the tree iteration of \mathbb{P}^J along the tree ${}^{<\omega}\omega_1$ (or the tree ${}^{<\omega}\omega$) the only L -generic filters for \mathbb{P}_n^J are the restrictions of G to level n nodes of tree. We proceed to construct a symmetric submodel of $L[G]$ which has the tree of \mathbb{P}_n^J -generic filters added by G but no branch through it. The symmetric model we construct satisfies AC_ω and the tree of \mathbb{P}_n^J -generic filters is Π_2^1 -definable in it. The reals of this model thus provide the desired β -model of Z_2 in which Σ_∞^1 -AC holds, but Π_2^1 -DC fails.

Our results also answer a long-standing open question of Zarach from [Zar96] about whether the Reflection Principle holds in models of ZFC^- . The *Reflection Principle* states that every formula can be reflected to a transitive set, and holds in ZFC by the Lévy-Montague reflection because every formula is reflected by some V_α . In the absence of the von Neumann hierarchy, it is not clear how to realize reflection, and indeed we show that it fails in $H_{\omega_1} \models ZFC^-$ of the symmetric model we construct.

2. JENSEN’S FORCING

Jensen’s poset \mathbb{P}^J , introduced in [Jen70], is constructed in L as the union of a continuous chain $\langle \mathbb{P}_\alpha \mid \alpha < \omega_1 \rangle$ of length ω_1 of countable perfect posets. At successor stages $\alpha + 1$, the principle \diamond is used to seal a certain countable collection of maximal antichains of \mathbb{P}_α , so that by the end of the construction every maximal antichain of \mathbb{P}^J is sealed and therefore countable.

Recall that a tree $T \subseteq {}^{<\omega}2$ is *perfect* if every node in T has a splitting node above it. Given two perfect trees T and S , let’s define their *meet* $T \wedge S$ as follows. Let $U_0 = T \cap S$, and, assuming U_α has been defined, let $U_{\alpha+1}$ be the set of all nodes in U_α which have a splitting node in U_α above them. At limit stages, take

intersections. Since U_0 is countable, there must be a countable stage α in the construction for which $U_\alpha = U_{\alpha+1}$. We define $T \wedge S = U_\alpha$. It is not difficult to see that either $T \wedge S = \emptyset$ or $T \wedge S$ is a perfect tree. In the latter case, it is the maximal perfect tree contained in $T \cap S$, that is every perfect tree $U \subseteq T \cap S$ is contained in $T \wedge S$. Recall the standard terminology that if $T \subseteq {}^{<\omega}2$ is a tree and s is a node in T , then T_s denotes the subtree of T consisting of all nodes compatible with s . The following proposition is straightforward to check.

Proposition 2.1. *If T, S , and R are perfect trees, then*

- (1) $(T \wedge S)_s = T_s \wedge S_s$ for every node $s \in T \wedge S$,
- (2) $(T \wedge S) \wedge R = T \wedge (S \wedge R)$ (\wedge is associative),
- (3) $(T \cup S) \wedge R = (T \wedge R) \cup (S \wedge R)$ (\wedge distributes over \cup).

Let's recall now some facts about perfect posets, which are subposets of Sacks forcing closed under certain basic operations.

Definition 2.2. We say that a collection \mathbb{P} of perfect trees ordered by inclusion is a *perfect poset*² if

- (1) $({}^{<\omega}2)_s \in \mathbb{P}$ for every $s \in {}^{<\omega}2$, and for every $T, S \in \mathbb{P}$,
- (2) $T \cup S \in \mathbb{P}$ (closed under unions),
- (3) $T \wedge S \in \mathbb{P}$, if $T \wedge S \neq \emptyset$ (closed under meets).

The smallest perfect poset, which we will denote by \mathbb{P}^{\min} , is the closure under finite unions of the collection $\{({}^{<\omega}2)_s \mid s \in {}^{<\omega}2\}$. Note that two perfect trees T and S are compatible under the inclusion ordering precisely when $T \wedge S \neq \emptyset$. So if $\mathbb{P} \subseteq \mathbb{Q}$ are perfect posets, then $T, S \in \mathbb{P}$ are compatible in \mathbb{Q} if and only if they are already compatible in \mathbb{P} . Standard arguments show that a generic filter G for a perfect poset \mathbb{P} is determined by the *generic real* r that is a branch through $T \in \mathbb{P}$ if and only if $T \in G$.

Given a perfect poset \mathbb{P} , we will denote by $\mathbb{P}^{<\omega}$ the ω -length finite-support product of \mathbb{P} . Conditions in $\mathbb{P}^{<\omega}$ are functions $p : \omega \rightarrow \mathbb{P}$ such that for all but finitely many n , $p(n) = {}^{<\omega}2$. We will sometimes abuse notation by writing $p = \langle T_0, \dots, T_{n-1} \rangle$ for conditions in $\mathbb{P}^{<\omega}$, meaning that all remaining coordinates are trivial.

Following [Abr84], we associate to a perfect poset \mathbb{P} , the poset $\mathbb{Q}(\mathbb{P})$ whose conditions are pairs (T, n) with $T \in \mathbb{P}$ and $n \in \omega$ ordered so that $(T_2, n_2) \leq (T_1, n_1)$ whenever $T_2 \subseteq T_1$, $n_2 \geq n_1$, and $T_1 \cap {}^{n_1}2 = T_2 \cap {}^{n_1}2$. We will refer to $\mathbb{Q}(\mathbb{P})$ as the *fusion poset* for \mathbb{P} because fusion arguments involving \mathbb{P} amount to producing a filter for $\mathbb{Q}(\mathbb{P})$ meeting sufficiently many dense sets. If $G \subseteq \mathbb{Q}(\mathbb{P})$ is V -generic, then the union of $T \cap {}^{n_1}2$ for $(T, n) \in G$ is the generic perfect tree \mathcal{T} added by $\mathbb{Q}(\mathbb{P})$. Note that $\mathcal{T} \leq T$ for every T that appears in some condition in G .

Let $\mathbb{Q}(\mathbb{P})^{<\omega}$ denote the ω -length finite-support product of the $\mathbb{Q}(\mathbb{P})$. Conditions in $\mathbb{Q}(\mathbb{P})^{<\omega}$ are function $q : \omega \rightarrow \mathbb{Q}(\mathbb{P})$ such that for all but finitely many n , $q(n) = ({}^{<\omega}2, 0)$, but we will sometimes abuse notation by writing the conditions as finite tuples.

²Note that this is not a standard definition. In the literature, a *perfect poset* is usually defined to be a collection \mathbb{P} of perfect trees such that ${}^{<\omega}2 \in \mathbb{P}$ and whenever $T \in \mathbb{P}$ and $s \in T$, then $T_s \in \mathbb{P}$.

Suppose that $G \subseteq \mathbb{Q}(\mathbb{P})^{<\omega}$ is V -generic. Let $\langle \mathcal{T}_n \mid n < \omega \rangle \in V[G]$ be the ω -length sequence of generic perfect trees derived from G and let $\dot{\mathcal{T}}_n$ be the canonical $\mathbb{Q}(\mathbb{P})^{<\omega}$ -names for the trees \mathcal{T}_n . In $V[G]$, let

$$\mathbb{U} = \{\mathcal{T}_n \wedge S \mid S \in \mathbb{P}, n \in \omega, \mathcal{T}_n \wedge S \neq \emptyset\}.$$

We will denote by \mathbb{P}^* the collection of perfect trees that is the closure under finite unions of \mathbb{P} and \mathbb{U} . We will argue that \mathbb{P}^* is a perfect poset extending \mathbb{P} . This will be the case even when G is only generic for a countable transitive model M containing \mathbb{P} . At successor stages $\alpha + 1$ in the construction of Jensen's poset, if \diamond codes a certain countable transitive model M with $\mathbb{P}_\alpha \in M$, we will extend \mathbb{P}_α to $\mathbb{P}_{\alpha+1} = \mathbb{P}_\alpha^*$ constructed in a generic extension $M[G]$ by $\mathbb{Q}(\mathbb{P}_\alpha)^{<\omega}$.

Suppose that M is a countable transitive model of $\text{ZFC}^- + \text{"}\mathcal{P}(\omega) \text{ exists"}$ and $\mathbb{P} \in M$ is a perfect poset. Clearly we have $\mathbb{Q}(\mathbb{P}) \in M$. We will argue that if $G \subseteq \mathbb{Q}(\mathbb{P})^{<\omega}$ is M -generic, then the poset \mathbb{P}^* constructed in $M[G]$ is a perfect poset with the property that every maximal antichain of \mathbb{P} that is an element of M remains maximal in \mathbb{P}^* . First, we need the following easy proposition.

Proposition 2.3. *Suppose \mathbb{P} is a perfect poset and a condition $q \in \mathbb{Q}(\mathbb{P})^{<\omega}$ forces that $\dot{\mathcal{T}}_n \wedge \dot{S} \neq \emptyset$ with $S \in \mathbb{P}$. Then there is a condition $\bar{q} \leq q$ with $\bar{q}(n) = (\bar{T}, m)$ and a node $s \in \bar{T} \cap {}^m 2$ for which $(\bar{T})_s \leq S$.*

Proof. Let $q(n) = (T, m)$. Since $q \Vdash \dot{\mathcal{T}}_n \wedge \dot{S} \neq \emptyset$, there must be some $s \in T \cap {}^m 2$ such that $U = T_s \wedge S \neq \emptyset$. Let \bar{T} be the perfect tree we get by replacing T_s with U in T and let $\bar{q} \leq q$ be such that $\bar{q}(n) = (\bar{T}, m)$ and $\bar{q}(i) = q(i)$ for all $i \neq n$. Note that the closure under unions property of perfect posets is needed to conclude that $\bar{T} \in \mathbb{P}$. \square

Proposition 2.4. *Suppose M is a countable transitive model of $\text{ZFC}^- + \text{"}\mathcal{P}(\omega) \text{ exists"}$ and $\mathbb{P} \in M$ is a perfect poset. If $G \subseteq \mathbb{Q}(\mathbb{P})^{<\omega}$ is M -generic and \mathbb{P}^* is constructed in $M[G]$ as above, then:*

- (1) \mathbb{P}^* is a perfect poset.
- (2) \mathbb{U} is dense in \mathbb{P}^* and $\{\mathcal{T}_n \mid n < \omega\}$ is a maximal antichain of \mathbb{P}^* .
- (3) Every maximal antichain of \mathbb{P} from M remains maximal in \mathbb{P}^* .

Proof. First, let's prove (2). We argue that every $T \in \mathbb{P}$ has some generic tree \mathcal{T}_n below it. Fix $T \in \mathbb{P}$. Let $q \in \mathbb{Q}(\mathbb{P})^{<\omega}$. Since q has finite support, we can choose some n such that $q(n) = ({}^{<\omega} 2, 0)$ and strengthen q to a condition \bar{q} such that $q(m) = \bar{q}(m)$ for all $m \neq n$ and $\bar{q}(n) = (T, 0)$. Clearly $\bar{q} \Vdash \dot{\mathcal{T}}_n \leq T$. So by density, there must be some such $\bar{q} \in G$.

Next, let's show that for $i \neq j$, $\mathcal{T}_i \wedge \mathcal{T}_j = \emptyset$, and so the generic trees \mathcal{T}_n form an antichain. Fix any condition $p \in \mathbb{Q}(\mathbb{P})^{<\omega}$ with $p(n) = (T_n, m_n)$. By strengthening further, we can assume that $m_i = m_j = m$. For each node s that is on level m of both T_i and T_j , we can choose disjoint perfect trees $U^{s,i} \subseteq (T_i)_s$ and $U^{s,j} \subseteq (T_j)_s$. Let S_i be the perfect tree which we get by replacing, for each s on level m of both T_i and T_j , $(T_i)_s$ with $U^{s,i}$ in T_i , and let S_j be obtained similarly. Now let $\bar{p} \leq p$ be the condition where on coordinate i , we put S_i instead of T_i and on coordinate j , we put S_j instead of T_j . Clearly \bar{p} forces that $\dot{\mathcal{T}}_i \wedge \dot{\mathcal{T}}_j = \emptyset$. It remains to show that every tree T in \mathbb{P}^* is compatible with some \mathcal{T}_n . If T is a finite union of trees one of which is in \mathbb{U} , then this is clear, and otherwise $T \in \mathbb{P}$, and we just argued that then some $\mathcal{T}_n \leq T$.

To prove (1), it suffices to show that \mathbb{P}^* is closed under meets. So suppose that $R = \bigcup_{i < n} R_i$ and $R' = \bigcup_{j < m} R'_j$, where $R_i, R'_j \in \mathbb{P} \cup \mathbb{U}$, and consider $R \wedge R' = \bigcup_{i < n, j < m} R_i \wedge R'_j$. We will argue that each $R_i \wedge R'_j$ is either empty or in $\mathbb{P} \cup \mathbb{U}$. If both R_i and R'_j are in \mathbb{P} , then the conclusion follows since \mathbb{P} is a perfect poset. If one of the R_i or R'_j is $\mathcal{T}_k \wedge S$ with $S \in \mathbb{P}$ and the other is $S' \in \mathbb{P}$, then $R_i \wedge R'_j = \mathcal{T}_k \wedge (S \wedge S')$ and the conclusion follows. Finally if $R_i = \mathcal{T}_k \wedge S$ and $R'_j = \mathcal{T}_{k'} \wedge S'$ with $S, S' \in \mathbb{P}$, then $R_i \wedge R'_j = (\mathcal{T}_k \wedge \mathcal{T}_{k'}) \wedge (S \wedge S')$. If $k = k'$, then the conclusion follows and if $k \neq k'$, we just argued above that $\mathcal{T}_k \wedge \mathcal{T}_{k'} = \emptyset$.

Let's now prove (3). Fix a maximal antichain $\mathcal{A} \in M$ of \mathbb{P} . It suffices to show that every tree in \mathbb{U} is compatible with some element of \mathcal{A} . So suppose that $\mathcal{T}_n \wedge S \in \mathbb{U}$ with $S \in \mathbb{P}$, and fix a condition $q \in G$ forcing that $\dot{\mathcal{T}}_n \wedge \dot{S} \neq \emptyset$. We will argue that there is a condition $q' \leq q$ with $q'(n) = (T', m)$ such that there is a node $s \in T' \cap {}^m 2$ and $A \in \mathcal{A}$ for which $T'_s \leq S, A$. By Proposition 2.3, there is $\bar{q} \leq q$ with $\bar{q}(n) = (\bar{T}, m)$ such that there is $s \in {}^m 2$ with $\bar{T}_s \leq S$. Since \mathcal{A} is maximal in \mathbb{P} , there is $A \in \mathcal{A}$ and $U \in \mathbb{P}$ such that $U \leq \bar{T}_s, A$. Let T' be the tree we get by replacing T'_s with U in \bar{T} and let $q' \leq q$ be the condition where $q'(k) = \bar{q}(k)$ for all $k \neq n$ and $q'(n) = (T', m)$. By density, there is some such condition $q' \in G$. But this means that $(\mathcal{T}_n)_s \leq \mathcal{T}_n \wedge S$ and $(\mathcal{T}_n)_s \leq A$, so that $\mathcal{T}_n \wedge S$ is compatible with $A \in \mathcal{A}$. \square

We let $\mathbb{U}^{<\omega}$ denote the collection of elements $p \in \mathbb{P}^{* <\omega}$ such that for all n , $p(n) \in \mathbb{U}$ or $p(n) = {}^{<\omega} 2$. Note that it follows from Proposition 2.4 that $\mathbb{U}^{<\omega}$ is dense in $\mathbb{P}^{* <\omega}$.

Proposition 2.5. *Suppose that M is a countable transitive model of $\text{ZFC}^- + \text{“}\mathcal{P}(\omega) \text{ exists”}$ and $\mathbb{P} \in M$ is a perfect poset. If $G \subseteq \mathbb{Q}(\mathbb{P})^{<\omega}$ is M -generic and \mathbb{P}^* is constructed in $M[G]$ as above, then every maximal antichain of $\mathbb{P}^{<\omega}$ from M remains maximal in $\mathbb{P}^{* <\omega}$.*

Proof. Fix a maximal antichain $\mathcal{A} \in M$ of $\mathbb{P}^{<\omega}$. It suffices to show that every condition $p \in \mathbb{U}^{<\omega}$ is compatible with some element of \mathcal{A} . Let

$$p = \langle \mathcal{T}_{i_0} \wedge S_0, \mathcal{T}_{i_1} \wedge S_1, \dots, \mathcal{T}_{i_m} \wedge S_m \rangle.$$

Note that the values i_n don't have to be distinct and we need to carefully address this possibility. Fix a condition $q \in G$ forcing that $\dot{\mathcal{T}}_{i_n} \wedge \dot{S}_n \neq \emptyset$ for every $n \leq m$. Repeatedly using the construction in the proof of Proposition 2.3 and going to a large enough level, we can find a condition $\bar{q} \leq q$ with $\bar{q}(n) = (\bar{T}_n, k_n)$ such that for every $n \leq m$, there is a node s_n on level k_{i_n} of \bar{T}_{i_n} such that $(\bar{T}_{i_n})_{s_n} \leq S_n$, and for $n \neq n'$, $s_n \neq s_{n'}$. Let $\bar{p} = \langle (\bar{T}_{i_0})_{s_0}, (\bar{T}_{i_1})_{s_1}, \dots, (\bar{T}_{i_m})_{s_m} \rangle$. Since \mathcal{A} is maximal in $\mathbb{P}^{<\omega}$ in \mathbb{P} , the condition \bar{p} is compatible with some $a \in \mathcal{A}$, and so for every $n \leq m$, we can let $U_n = \bar{p}(n) \wedge a(n) \in \mathbb{P}$. Now, for every $i \in \{i_n \mid n \leq m\}$, let R_i be the tree we get by replacing $(\bar{T}_i)_{s_n}$ with U_n in \bar{T}_i whenever $i_n = i$. Let r be the condition such that $r(i) = (R_i, k_i)$ for $i \in \{i_n \mid n \leq m\}$ and $r(i) = \bar{q}(i)$ otherwise. By density, there is some such $r \in G$. But this means that $(\mathcal{T}_{i_n})_{s_n} \leq S_n \wedge a(n)$, so that p is compatible with a . \square

All our constructions will take place inside the constructible universe L . For reasons that will become obvious during the course of the constructions, we will restrict ourselves to countable models of $\text{ZFC}^- + \text{“}\mathcal{P}(\omega) \text{ exists”}$ which happen to be initial segments L_α of L . So let's call countable L_α satisfying $\text{ZFC}^- + \text{“}\mathcal{P}(\omega) \text{ exists”}$

suitable models. Relevant examples of suitable models for us will be transitive collapses of countable $M \prec L_{\omega_2}$.

We are now ready to review the construction of Jensen's perfect poset \mathbb{P}^J , which will use the \diamond -principle to anticipate and seal maximal antichains. So let's start by fixing a canonically defined \diamond -sequence $\langle S_\alpha \mid \alpha < \omega_1 \rangle$. Note that if M is a suitable model and $\delta = \omega_1^M$, then $\langle S_\alpha \mid \alpha < \delta \rangle$ is an element of M .

Jensen's poset \mathbb{P}^J will be the union of the following increasing sequence $\langle \mathbb{P}_\alpha \mid \alpha < \omega_1 \rangle$ of perfect posets. Let $\mathbb{P}_0 = \mathbb{P}^{\min}$. At limit stages, we will take unions. Suppose \mathbb{P}_α has been defined. We let $\mathbb{P}_{\alpha+1} = \mathbb{P}_\alpha$, unless the following happens. Suppose S_α codes a well-founded and extensional binary relation $E \subseteq \alpha \times \alpha$ such that the collapse of E is a suitable model M_α with $\mathbb{P}_\alpha \in M_\alpha$ and $\alpha = \omega_1^{M_\alpha}$. In this case, we take the L -least M_α -generic filter $G \subseteq \mathbb{Q}(\mathbb{P}_\alpha)^{<\omega}$ and let $\mathbb{P}_{\alpha+1} = \mathbb{P}_\alpha^*$ as constructed in $M_\alpha[G]$.

As we observed in Proposition 2.4, $\mathbb{P}_{\alpha+1}$ is a perfect poset with the property that every maximal antichain of \mathbb{P}_α in M_α remains maximal in $\mathbb{P}_{\alpha+1}$. Also, by Proposition 2.5, every maximal antichain of $\mathbb{P}_\alpha^{<\omega}$ in M_α remains maximal in $\mathbb{P}_{\alpha+1}^{<\omega}$. Now let's argue that every maximal antichain of $\mathbb{P}_\alpha^{<\omega}$ in M_α remains maximal in the final poset $\mathbb{P}^{<\omega}$. It suffices to argue that the models M_α form an increasing sequence, which follows because if $\beta > \alpha$, then $\beta = \omega_1^{M_\beta}$, and therefore M_β has S_α as an element and can collapse it to obtain M_α . This shows that every maximal antichain of $\mathbb{P}_\alpha^{<\omega}$ that is an element of M_α is sealed.

Theorem 2.6. *The finite-support product $\mathbb{P}^{J<\omega}$ of Jensen's poset \mathbb{P}^J has the ccc.*

Proof. Fix a maximal antichain \mathcal{A} of $\mathbb{P}^{J<\omega}$. Choose some transitive $M \prec L_{\omega_2}$ of size ω_1 with $\mathcal{A} \in M$. We can decompose M as the union of a continuous elementary chain of countable substructures

$$X_0 \prec X_1 \prec \cdots \prec X_\alpha \prec \cdots \prec M$$

with $\mathcal{A} \in X_0$. By properties of \diamond , there is some α such that $\alpha = \omega_1 \cap X_\alpha$, $\mathbb{P}_\alpha = \mathbb{P}^J \cap X_\alpha$, and S_α codes X_α . Let M_α be the transitive collapse of X_α . Then \mathbb{P}_α is the image of \mathbb{P}^J under the collapse and α is the image of ω_1 . Let $\bar{\mathcal{A}} = \mathcal{A} \cap X_\alpha$ be the image of \mathcal{A} under the collapse. So at stage α in the construction of \mathbb{P}^J , we chose a forcing extension $M_\alpha[G]$ of M_α by $\mathbb{Q}(\mathbb{P}_\alpha)^{<\omega}$ and let $\mathbb{P}_{\alpha+1} = \mathbb{P}_\alpha^*$ as constructed in $M_\alpha[G]$. Thus, by our observation above, $\bar{\mathcal{A}}$ remains maximal in $\mathbb{P}^{J<\omega}$, and hence $\bar{\mathcal{A}} = \mathcal{A}$ is countable. \square

Finally, we would like to observe that every maximal antichain of $\mathbb{P}_{\alpha+1}$ from $M_\alpha[G]$, in particular the antichain $\langle \mathcal{T}_n \mid n < \omega \rangle$ of generic perfect trees, remains maximal in \mathbb{P}^J .

Proposition 2.7. *Suppose that $\mathcal{A} \in M_\alpha[G]$ is a maximal antichain of $\mathbb{P}_{\alpha+1}$ (or $\mathbb{P}_{\alpha+1}^{<\omega}$). Then \mathcal{A} remains maximal in \mathbb{P}^J (or $\mathbb{P}^{J<\omega}$).*

Proof. Fix $\beta > \alpha$. We already argued that $M_\alpha \in M_\beta$. Since M_β sees that M_α is countable, it has some M -generic filter for $\mathbb{Q}(\mathbb{P}_\alpha)^{<\omega}$. But since G_α was chosen to be the L -least such filter, then by suitability, M_β must contain G_α . So $M_\alpha[G_\alpha] \subseteq M_\beta$. \square

3. THE KANOVEI-LYUBETSKY THEOREM

We will reprove here the Kanovei-Lyubetsky theorem from [KL] showing that Jensen's poset \mathbb{P}^J from Section 2, has the property that in a forcing extension $L[G]$

of L by the finite support ω -length product $\mathbb{P}^{J<\omega}$, the only L -generic reals for \mathbb{P} are the ω -many slices of G . This is a generalization of Jensen's uniqueness of generic filters property to products.

Suppose that \mathbb{P} is a perfect poset and G is a generic filter for the product $\mathbb{P}^{<\omega}$. We will call x_n the real on the n -th coordinate of G and let \dot{x}_n be its canonical name.

For the next lemma, we suppose that \mathbb{P} is a countable perfect poset that is an element of a suitable model M . We should think of \mathbb{P} as one of the perfect posets \mathbb{P}_α arising at stage α in the construction of Jensen's poset \mathbb{P}^J and of M as the model M_α from that stage.

Theorem 3.1 (Kanovei-Lyubetsky, [KL]). *In M , suppose that \dot{r} is a $\mathbb{P}^{<\omega}$ -name for a real such that for all $n \in \omega$, $\mathbb{1} \Vdash_{\mathbb{P}^{<\omega}} \dot{r} \neq \dot{x}_n$. Then in a forcing extension $M[G]$ by $\mathbb{Q}(\mathbb{P})^{<\omega}$, for every generic perfect tree \mathcal{T}_n , conditions forcing that $\dot{r} \notin [\mathcal{T}_n]$ are dense in $\mathbb{P}^{*<\omega}$.*

Proof. Fix a condition $p \in \mathbb{P}^{*<\omega}$ and $d \in \omega$. We need to find $p' \leq p$ such that $p' \Vdash \dot{r} \notin [\mathcal{T}_d]$. Since $\mathbb{U}^{<\omega}$ is dense in $\mathbb{P}^{*<\omega}$, we can assume without loss that $p \in \mathbb{U}^{<\omega}$. So let

$$p = \langle \mathcal{T}_{i_0} \wedge S_0, \mathcal{T}_{i_1} \wedge S_1, \dots, \mathcal{T}_{i_m} \wedge S_m \rangle.$$

By strengthening if necessary, we can assume that for some $n \leq m$, $i_n = d$. As we argued in the proof of Proposition 2.5, there is a condition $q \in G$ with $q(n) = (T_n, k_n)$ such that for every $n \leq m$, there is a node s_n on level k_{i_n} of T_{i_n} such that $(T_{i_n})_{s_n} \leq S_n$ and for $n \neq n'$, $s_n \neq s_{n'}$.

Now we are going to construct a condition $a_q = \langle W_0, \dots, W_{l'}, \dots, W_l \rangle \in \mathbb{P}^{<\omega}$ associated to q and satisfying the following properties.

- (1) For every $n \leq m \leq l'$, $W_n \leq (T_{i_n})_{s_n}$.
- (2) For every node s on level k_d of T_d , there is $i \leq l'$ such that $W_i \leq (T_d)_s$.
- (3) For every $n \leq l'$, $a_q \Vdash_{\mathbb{P}^{<\omega}} \dot{r} \notin [W_n]$.

Let $a = \langle (T_{i_0})_{s_0}, (T_{i_1})_{s_1}, \dots, (T_{i_m})_{s_m}, (T_d)_{r_0}, \dots, (T_d)_{r_k} \rangle$, where r_0, \dots, r_k are nodes on level k_d of T_d excluding those amongst the s_n . Since $\mathbb{1}_{\mathbb{P}^{<\omega}} \Vdash \dot{r} \neq \dot{x}_n$ for all $n \in \omega$, we can strengthen a to a condition $a_q = \langle W_0, W_1, \dots, W_{m+k+2}, \dots, W_l \rangle$ forcing that $\dot{r} \notin [W_n]$ for all $n \leq m+k+2$.

Using a_q , we construct the following condition $\bar{q} \leq q$. For $i \in \{i_n \mid n \leq m\}$, let R_i be the tree we get by replacing $(T_i)_{s_n}$ with W_n in T_i whenever $i_n = i$ and if $i = d$, then we also replace all T_{r_n} with the appropriate W_j as well. Let $\bar{q}(i) = (R_i, k_i)$ for $i \in \{i_n \mid n \leq m\}$ and $\bar{q}(i) = q(i)$ otherwise. By density, some such condition \bar{q} , constructed from a_q , is in G . It follows that each $(T_{i_n})_{s_n} \leq W_n \wedge S_n$ and \mathcal{T}_d has a level k_d such that for every node s on level k_d , $\mathcal{T}_d \leq W_j$ for some j . The first part gives us that $p = \langle \mathcal{T}_{i_0} \wedge S_0, \mathcal{T}_{i_1} \wedge S_1, \dots, \mathcal{T}_{i_m} \wedge S_m \rangle$ is compatible with a_q . Finally, we would like to argue that $a_q \Vdash_{\mathbb{P}^{*<\omega}} \dot{r} \notin [\mathcal{T}_d]$.

Let H^* be any V -generic filter for $\mathbb{P}^{*<\omega}$ containing a_q ($V = M[G]$). Since every maximal antichain of $\mathbb{P}^{<\omega}$ in M remains maximal in $\mathbb{P}^{*<\omega}$, it follows that H^* restricts to an M -generic filter H for $\mathbb{P}^{<\omega}$ and $a_q \in H$. Thus in $M[H]$, \dot{r}_H is not a branch through any W_n . But this is absolute, and so \dot{r}_H is not a branch through \mathcal{T}_d in $V[H^*]$. \square

Theorem 3.2 (Kanovei-Lyubetsky, [KL]). *Suppose $H \subseteq \mathbb{P}^{J<\omega}$ is L -generic. If $r \in L[H]$ is L -generic for \mathbb{P}^J , then $r = x_n$ for some $n < \omega$.*

Proof. Let's suppose that $r \in L[H]$ is a real which is not one of the x_n . Let \dot{r} be a nice $\mathbb{P}^{J<\omega}$ -name for r such that for all $n \in \omega$, $\mathbb{1}_{\mathbb{P}^{J<\omega}} \Vdash \dot{r} \neq \dot{x}_n$.

Choose some transitive model $M \prec L_{\omega_2}$ of size ω_1 with $\dot{r} \in N$. We can decompose M as the union of a continuous elementary chain of countable substructures

$$X_0 \prec X_1 \prec \cdots \prec X_\alpha \prec \cdots \prec M$$

with $\dot{r} \in X_0$. By properties of \diamond , there is some α such that $\alpha = \omega_1 \cap X_\alpha$, $\mathbb{P}_\alpha = \mathbb{P}^J \cap X_\alpha$, and S_α codes X_α . Let M_α be the collapse of X_α . Then \mathbb{P}_α is the image of \mathbb{P}^J under the collapse and α is the collapse of ω_1 . Clearly \dot{r} is fixed by the collapse because it is a nice name and all antichains of $\mathbb{P}^{J<\omega}$ are countable (by Theorem 2.6). So at stage α in the construction of \mathbb{P}^J , we chose a forcing extension $M_\alpha[G]$ by $\mathbb{Q}(\mathbb{P}_\alpha)^{<\omega}$ and let $\mathbb{P}_{\alpha+1} = \mathbb{P}_\alpha^*$ as constructed in $M_\alpha[G]$. By elementarity, M_α satisfies that $\mathbb{1}_{\mathbb{P}_\alpha^{<\omega}} \Vdash \dot{r} \neq \dot{x}_n$ for all $n \in \omega$. Thus, by Theorem 3.1, for every $n < \omega$, $\mathbb{P}_{\alpha+1}^{<\omega}$ has a maximal antichain $\mathcal{A}_n \in M_\alpha[G]$ consisting of conditions q such that $q \Vdash_{\mathbb{P}_{\alpha+1}^{<\omega}} \dot{r} \notin [\mathcal{T}_n]$. It follows, using Proposition 2.7, that all the antichains \mathcal{A}_n remain maximal in $\mathbb{P}^{J<\omega}$. Also, by Proposition 2.7, $\langle \mathcal{T}_n \mid n < \omega \rangle$ remains maximal in \mathbb{P}^J .

So let's argue that if $q \in \mathcal{A}_n$, then $q \Vdash_{\mathbb{P}^{J<\omega}} \dot{r} \notin [\mathcal{T}_n]$. Let $\bar{H}^* \subseteq \mathbb{P}^{J<\omega}$ be an L -generic filter containing q and let \bar{H} be the restriction of \bar{H}^* to $\mathbb{P}_{\alpha+1}^{<\omega}$. Since $q \in \bar{H}$, it follows that $\dot{r}_{\bar{H}} \notin [\mathcal{T}_n]$ holds in $M_\alpha[G_\alpha][\bar{H}]$, but this statement is absolute and so also holds in $L[\bar{H}^*]$. Since H must meet every \mathcal{A}_n , it holds in $L[H]$ that $\dot{r}_H = r$ is not a branch through any \mathcal{T}_n . So r cannot be L -generic for \mathbb{P}^J . \square

4. FINITE ITERATIONS OF PERFECT POSETS

An iteration of perfect posets is an iteration of forcing notions in which every initial segment forces that the next poset is perfect. Here we will only be dealing with finite iterations of perfect posets, so we are not concerned with issues of support.

Definition 4.1. A *finite iteration of perfect posets* is a finite iteration

$$\mathbb{P}_n = \mathbb{Q}_0 * \dot{\mathbb{Q}}_1 * \cdots * \dot{\mathbb{Q}}_{n-1}$$

such that \mathbb{Q}_0 is a perfect poset and for $1 \leq i < n$,

$$\mathbb{1}_{\mathbb{P}_i} \Vdash \text{“}\dot{\mathbb{Q}}_i \text{ is a perfect poset”}.$$

Suppose $G \subseteq \mathbb{P}_n$ is V -generic. For $1 \leq i < n$, let G_i be the restriction of G to \mathbb{P}_i . Let $G(0) = G_1$ and for $1 \leq i < n$, let $G(i) = \{p(i)_{G_i} \mid p \in G\}$. Let r_i be the unique real determined by $G(i)$. It is not difficult to see that the sequence of reals $\vec{r} = \langle r_0, \dots, r_{n-1} \rangle$ determines G . Elements of G_1 are trees with r_0 as a branch, and inductively, elements of G_{i+1} are conditions $p \in \mathbb{P}_i$ such that $p \restriction i \in G_i$ and $r_i \in p(i)_{G_i}$.

The analogue of the fusion poset $\mathbb{Q}(\mathbb{P})$ for a finite iteration \mathbb{P}_n of perfect posets is the fusion poset $\mathbb{Q}(\mathbb{P}_n)$ whose conditions are pairs (p, F) with $p \in \mathbb{P}_n$ and $F : n \rightarrow \omega$, ordered so that $(p_2, F_2) \leq (p_1, F_1)$ whenever $p_2 \leq p_1$ and for every $i < n$, we have $F_2(i) \geq F_1(i)$ and

$$p_2 \restriction i \Vdash p_1(i) \cap^{F_1(i)} 2 = p_2(i) \cap^{F_1(i)} 2.$$

Fusion arguments with names for perfect trees require that we have some information about a fixed level n of the tree. We will now argue that there are densely many conditions in $\mathbb{Q}(\mathbb{P}_n)$ where this is the case.

Suppose $p \in \mathbb{P}_n$ and $\sigma : n \rightarrow {}^{<\omega}2$. Following [Abr84], let's define, by induction on n , what it means for σ to *lie* on p .³ For $n = 1$, we shall say that σ *lies* on p whenever $\sigma(0) \in p(0)$. If σ lies on p , we shall denote by $p \mid \sigma$ the condition $p(0)_{\sigma(0)}$. Note that $p \mid \sigma \leq p$. So suppose that we have defined when σ lies on p for $p \in \mathbb{P}_n$, and for σ which lies on p , we have defined $p \mid \sigma$ so that $p \mid \sigma \leq p$. Let $p \in \mathbb{P}_{n+1}$. We define that σ lies on p if $\sigma \upharpoonright n$ lies on $p \upharpoonright n$ and $(p \upharpoonright n) \mid (\sigma \upharpoonright n) \Vdash \sigma(n) \in p(n)$. If σ lies on p , we shall denote by $p \mid \sigma$ the condition \bar{p} such that $\bar{p} \upharpoonright n = (p \upharpoonright n) \mid (\sigma \upharpoonright n)$ and $\bar{p}(n) = \dot{T}$, where \dot{T} is a \mathbb{P}_n -name that is interpreted as $p(n)_{\sigma(n)}$ by any \mathbb{P}_n -generic filter containing $(p \upharpoonright n) \mid (\sigma \upharpoonright n)$ and as $p(n)$ otherwise. Clearly this gives that $p \mid \sigma \leq p$.

Definition 4.2. Let $F : n \rightarrow \omega$ and $\sigma : n \rightarrow {}^{<\omega}2$. We shall say that σ *lies on levels* F if $\sigma(i) \in {}^{F(i)}2$ for all $i < n$.⁴ We shall say that a pair (p, F) with $p \in \mathbb{P}_n$ is *determined* if for every σ lying on levels F , either σ lies on p or there is some $i < n$ such that $\sigma \upharpoonright i$ lies on $p \upharpoonright i$ and $(p \upharpoonright i) \mid (\sigma \upharpoonright i) \Vdash \sigma(i) \notin p(i)$. If σ lies on levels F and lies on p , we shall say that σ *lies on* (p, F) .

Observe that whenever a pair (p, F) is determined, for every σ which lies on p , $(p \mid \sigma) \upharpoonright i$ decides the $F(i)$ -th level of $p(i)$, and indeed $(p \mid \sigma) \upharpoonright i$ forces that $p(i)$ is the union of the $p \mid \sigma'(i)$ for $\sigma' \upharpoonright i = \sigma \upharpoonright i$ that lie on (p, F) .

Proposition 4.3. *Suppose \mathbb{P}_n is a finite iteration of perfect posets.*

- (1) *Determined conditions (p, F) are dense in $\mathbb{Q}(\mathbb{P}_n)$.*
- (2) *Given a determined condition (p, F) , the finite set*

$$\{p \mid \sigma \mid \sigma \text{ lies on } (p, F)\}$$

is a maximal antichain below p in \mathbb{P}_n .

Proposition 4.4. *Suppose \mathbb{P}_n is a finite iteration of perfect posets and $D \subseteq \mathbb{P}_n$ is open dense. For any determined $(p, F) \in \mathbb{Q}(\mathbb{P}_n)$, there is a determined $(q, F) \leq (p, F)$ satisfying*

$$q \mid \sigma \in D \text{ whenever } \sigma \text{ lies on } (q, F).$$

For details of proofs, see [Abr84].

Since we will be working mainly with determined conditions, we will now introduce a kind of normal form for them.

Definition 4.5. Suppose \mathbb{P}_n is a finite iteration of perfect posets and $\sigma : n \rightarrow {}^{<\omega}2$. Let's call a condition $p \in \mathbb{P}_n$ a σ -*condition* if

- (1) $p(0) \leq ({}^{<\omega}2)_{\sigma(0)}$,
- (2) for all $1 \leq i < n$, $p \upharpoonright i \Vdash p(i) \leq ({}^{<\omega}2)_{\sigma(i)}$.

Suppose that X_F is a collection of $\sigma : n \rightarrow {}^{<\omega}2$ lying on levels $F : n \rightarrow \omega$. An X_F -*assignment* is a function $\varphi : X_F \rightarrow \mathbb{P}_n$ such that each $\varphi(\sigma)$ is a σ -condition and $\varphi(\sigma) \upharpoonright i = \varphi(\sigma') \upharpoonright i$ whenever $\sigma \upharpoonright i = \sigma' \upharpoonright i$.

To motivate these definitions, consider a determined condition $(p, F) \in \mathbb{Q}(\mathbb{P}_n)$ and let X_F^p be the set of all σ that lie on it. In this case, the map φ_p defined by $\varphi_p(\sigma) = p \mid \sigma$ for all $\sigma \in X_F^p$ is clearly an X_F^p -assignment. Thus, a determined condition (p, F) gives us a natural X_F -assignment, and now conversely we would like

³Abraham [Abr84] uses the terminology σ is *consistent* with p .

⁴Abraham [Abr84] uses the terminology σ is *bounded* by F .

to argue that any X_F -assignment has a naturally associated determined condition (q, F) .

Suppose X_F and φ are as in Definition 4.5. Observe that given any $\sigma, \sigma' \in X_F$, either $\sigma(0) = \sigma'(0)$ and so $\varphi(\sigma)(0) = \varphi(\sigma')(0)$, or $\sigma(0) = s \neq t = \sigma'(0)$ are two nodes on level $F(0)$ such that $\varphi(\sigma)(0) \leq (\omega 2)_s$ and $\varphi(\sigma')(0) \leq (\omega 2)_t$. More generally, if $\sigma \neq \sigma'$, then there is some least i such that $\sigma \upharpoonright i = \sigma' \upharpoonright i$ and there are nodes $\sigma(i) = s \neq t = \sigma'(i)$ on level $F(i)$ such that $\varphi(\sigma) \upharpoonright i \Vdash \varphi(\sigma)(i) \leq (\omega 2)_s$ and $\varphi(\sigma') \upharpoonright i \Vdash \varphi(\sigma')(i) \leq (\omega 2)_t$. It follows, in particular, that for any $i < n$, the conditions $\varphi(\sigma) \upharpoonright i$ for $\sigma \in X_F$ form an antichain.

Proposition 4.6. *Suppose X_F is a collection of $\sigma : n \rightarrow \omega 2$ on level some $F : n \rightarrow \omega$ and φ is an X_F -assignment. Then there is a determined condition $(q, F) \in \mathbb{Q}(\mathbb{P}_n)$ such that*

- (1) σ lies on (q, F) if and only if $\sigma \in X_F$,
- (2) for every $\sigma \in X_F$,

$$q \upharpoonright \sigma(0) = \varphi(\sigma)(0),$$

and for all $i < n$, $(q \upharpoonright \sigma) \upharpoonright i$ and $\varphi(\sigma) \upharpoonright i$ extend each other and force

$$q \upharpoonright \sigma(i) = \varphi(\sigma)(i).$$

Proof. Let $q(0)$ be the union of the $\varphi(\sigma)(0)$ for $\sigma \in X_F$, which is in the perfect poset \mathbb{Q}_0 . Let $q(1)$ be the (canonical) \mathbb{P}_1 -name for the tree which is the union of the collection of trees given by the interpretation of the name

$$\{(\varphi(\sigma)(1), \varphi(\sigma)(0)) \mid \sigma \in X_F\}.$$

Since each $\varphi(\sigma)(0) \Vdash \varphi(\sigma)(1) \in \mathbb{Q}_1$, and \mathbb{Q}_1 is forced to be a perfect poset, it follows that $q(0) \Vdash q(1) \in \mathbb{Q}_1$. Let's see what $q(1)_G$ looks like in a forcing extension $V[G]$ by \mathbb{P}_1 . If $\varphi(\sigma)(0)$ and $\varphi(\sigma')(0)$ are in G , then $\sigma(0) = \sigma'(0) = s$ for some s , and so $\varphi(\sigma)(0) = \varphi(\sigma')(0) = p$ for some p . Thus, the interpretation $q(1)_G$ is the tree which is the union of the $\varphi(\sigma)(1)_G$ for $\sigma(0) = s$. Similarly, let $q(i)$ be the \mathbb{P}_i -name for the tree which is the union of the collection of trees given by the interpretation of the name $\{(\varphi(\sigma)(i), \varphi(\sigma) \upharpoonright i) \mid \sigma \in X_F\}$. Again, we have $q \upharpoonright i \Vdash q(i) \in \mathbb{Q}_i$. Let's see now what $q(i)_G$ looks like in a forcing extension $V[G]$ by \mathbb{P}_i . Since $\varphi(\sigma) \upharpoonright i$ for $\sigma \in X_F$ form an antichain, if $\varphi(\sigma) \upharpoonright i$ and $\varphi(\sigma') \upharpoonright i$ are both in G , then $\sigma' \upharpoonright i = \sigma \upharpoonright i = \tau$ for some τ . So the interpretation $q(i)_G$ is the union of the $\varphi(\sigma)(i)_G$ for $\sigma \upharpoonright i = \tau$.

First, we argue that every $\sigma \in X_F$ lies on (q, F) and simultaneously show (2). So fix some $\sigma \in X_F$. By construction $\sigma(0) \in q(0)$ and $q(0)_{\sigma(0)} = \varphi(\sigma)(0)$. So assume inductively that for some $i < n$,

- (1) $\sigma \upharpoonright i$ lies on $q \upharpoonright i$,
- (2) $(q \upharpoonright i) \upharpoonright (\sigma \upharpoonright i) \leq \varphi(\sigma) \upharpoonright i$,
- (3) $\varphi(\sigma) \upharpoonright i \leq (q \upharpoonright i) \upharpoonright (\sigma \upharpoonright i)$.

Suppose that $G \subseteq \mathbb{P}_i$ is a V -generic filter containing $(q \upharpoonright i) \upharpoonright (\sigma \upharpoonright i)$. Then also, by assumption, $\varphi(\sigma) \upharpoonright i \in G$. By definition of q , we have

$$(q(i)_G)_{\sigma(i)} = \varphi(\sigma)(i)_G.$$

So $\sigma \upharpoonright i + 1$ lies on $q \upharpoonright i + 1$ and we have

$$(q \upharpoonright i + 1) \upharpoonright (\sigma \upharpoonright i + 1) \leq \varphi(\sigma) \upharpoonright i + 1 \text{ and } \varphi(\sigma) \upharpoonright i + 1 \leq (q \upharpoonright i + 1) \upharpoonright (\sigma \upharpoonright i + 1).$$

Now suppose that $\tau : n \rightarrow {}^{<\omega}2$ lies on (q, F) . By definition of q , it follows that $\tau(0) = \sigma(0)$ for some $\sigma \in X_F$. So suppose inductively that for some $i < n$, there is $\sigma \in X_F$ such that $\tau \upharpoonright i = \sigma \upharpoonright i$. Since τ lies on q , it follows that $(q \upharpoonright i) \mid (\sigma \upharpoonright i) \Vdash \tau(i) \in q(i)$. But then there is some σ' such that $\sigma \upharpoonright i = \sigma' \upharpoonright i$ and $\tau(i) = \sigma'(i)$. So in the last step, we will obtain $\sigma \in X_F$ such that $\sigma = \tau$. \square

We shall call the condition q constructed in Proposition 4.6 the *amalgamation* of φ .

Proposition 4.7. *Suppose (p, F) is determined and q is the amalgamation of the X_F^p -assignment φ_p . Then $q \leq p$ and $p \leq q$.*

Proof. Clearly $p(0) = q(0)$. So let's suppose inductively that for some $i < n$, we have $q \upharpoonright i \leq p \upharpoonright i$ and $p \upharpoonright i \leq q \upharpoonright i$. We will argue that both $p \upharpoonright i$ and $q \upharpoonright i$ force $p(i) = q(i)$. So suppose $G \subseteq \mathbb{P}_i$ is a V -generic filter containing $p \upharpoonright i$, and hence also $q \upharpoonright i$. Since the conditions $(p \mid \sigma) \upharpoonright i$ for $\sigma \in X_F^p$ form a maximal antichain below $p \upharpoonright i$, it follows that one such $(p \mid \sigma) \upharpoonright i$ is in G . Thus, $p(i)_G$ is the union of the $(p \mid \sigma'(i))_G$ for $\sigma' \in X_F^p$ with $\sigma' \upharpoonright i = \sigma \upharpoonright i$. Similarly, there is some $\tau \in X_F^p$ such that $(q \mid \tau) \upharpoonright i \in G$, so we must have $\tau \upharpoonright i = \sigma \upharpoonright i$. Also, $q(i)_G$ is the union of the $(q \mid \sigma'(i))_G$ for $\sigma' \in X_F^p$ with $\sigma' \upharpoonright i = \sigma \upharpoonright i$. Now, by Proposition 4.6, $(p \mid \sigma) \upharpoonright i \Vdash q \mid \sigma'(i) = p \mid \sigma'(i)$ for all $\sigma' \in X_F^p$ with $\sigma' \upharpoonright i = \sigma \upharpoonright i$. So $p(i)_G = q(i)_G$. \square

Let us say that a map $\tau : n \rightarrow {}^{<\omega}2$ extends another map $\sigma : n \rightarrow {}^{<\omega}2$ if $\tau(i)$ extends $\sigma(i)$ for every $i < n$. Now we would like to argue that given a determined condition (q, F) and strengthenings $q_\sigma \leq q \mid \sigma$ for every σ lying on (q, F) , we can obtain a determined condition $(\bar{q}, \bar{F}) \leq (q, F)$ such that for every σ lying on (q, F) , there is some τ lying on (\bar{q}, \bar{F}) extending σ with $\bar{q} \upharpoonright \tau \leq q_\sigma$. Indeed, we will get the following stronger statement.

Proposition 4.8. *Suppose $(q, F) \in \mathbb{Q}(\mathbb{P}_n)$ is determined and for every σ which lies on (q, F) , there is a finite set \mathcal{X}_σ of conditions $p \leq q \mid \sigma$. Then there is a condition $(\bar{q}, \bar{F}) \leq (q, F)$, also determined, such that for every σ which lies on (q, F) and $p \in \mathcal{X}_\sigma$, there is τ which lies on (\bar{q}, \bar{F}) and extends σ having $\bar{q} \upharpoonright \tau \leq p$. Moreover, for every τ which lies on (\bar{q}, \bar{F}) , $\bar{q} \upharpoonright \tau \leq p$ for some $p \in \mathcal{X}_\sigma$ with τ extending σ .*

Proof. First, suppose $n = 1$. Let (T, n) be a condition in $\mathbb{Q}(\mathbb{P})$ and suppose that for every node t on level n of T , we have a finite set \mathcal{X}_t of trees $S \leq T_t$. Let T' be the tree we get by replacing T_t with the union of S in \mathcal{X}_t for every t on level n of T . Fix t on level n of T' . Find a large enough level $n_t > n$ such that for every s on level n_t of T' , we can thin out T'_s to P^s with the property that $P^s \leq S$ for some $S \in \mathcal{X}_t$, and moreover for every $S \in \mathcal{X}_t$, there is some node s such that $P^s \leq S$. Let \bar{T} be the tree we obtain by replacing T'_s with P^s in T' . Let $\bar{n} > n_t$ for every t on level n of T' . Clearly the condition (\bar{T}, \bar{n}) has all the desired properties.

Next, let's consider the case $n = 2$. Let (q, F) be a condition in $\mathbb{Q}(\mathbb{P}_2)$ and suppose that for every σ which lies on (q, F) , we have a finite set \mathcal{X}_σ of conditions $p \leq q \mid \sigma$. For every node t on level $F(0)$ of $q(0)$, let \mathcal{X}_t be the set of all trees $p(0)$ with $p \in \mathcal{X}_\sigma$ for $\sigma(0) = t$. Using the case $n = 1$, let (\bar{T}, \bar{n}) be the condition for the sets \mathcal{X}_t . Fix a node s on level \bar{n} of \bar{T} and let t_s be the node on level $F(0)$ of \bar{T} which s extends. Since (q, F) was determined, \bar{T}_s decides the $F(1)$ -th level of $q(1)$. So let \bar{T}^s be a \mathbb{P}_1 -name for the tree we get by replacing, for every node t on level $F(1)$ of $q(1)$, each $q(1)_t$ with the union of $p(1)$ for $p \in \mathcal{X}_\sigma$ with $\sigma = \{(0, t_s), (1, t)\}$

and $\bar{T}_s \leq p(0)$. Strengthen \bar{T}_s to P^s deciding a level $m_s > F(1)$ such that for every node u on level m_s of \bar{T}^s above a node t on level $F(1)$, we can thin out \bar{T}_u^s to $\dot{P}^{s,u}$ such that $\dot{P}^{s,u} \leq p(1)$ for some $p \in \mathcal{X}_\sigma$ with $\sigma = \{(0, t_s), (1, t)\}$ and $\bar{T}_s \leq p(0)$, and moreover for every such $p(1)$, there is some node u such that $\dot{P}^{s,u} \leq p(1)$. We can assume, with some more thinning out, that all $m_s = m$ for a fixed m . Let \bar{T}' be the tree we get by replacing each \bar{T}_s with P^s in \bar{T} . Let \dot{T}'^s be a \mathbb{P}_1 -name for the tree we get by replacing, for every node u on level m of \dot{T}^s , each \dot{T}_u^s with $\dot{P}^{s,u}$. Let $\bar{F} = \{(0, \bar{n}), (1, m)\}$. Let $X_{\bar{F}}$ consist of $\sigma : 2 \rightarrow {}^{<\omega}2$ such that $\sigma(0) = s$ on level \bar{n} of \bar{T}' and $\sigma(1) = u$ on level m of \dot{T}'^s . Let φ be the $X_{\bar{F}}$ -assignment such that $\varphi(\sigma) = \{(0, \bar{T}'_{\sigma(0)}), (1, \dot{T}'_{\sigma(1)}{}^{(0)})\}$. Let \bar{q} be the amalgamation of φ . Clearly the condition (\bar{q}, \bar{F}) has all the desired properties.

Finally, assuming that the statement holds for n , let's argue that it holds for $n + 1$ by mimicking how the argument moves from $n = 1$ to $n = 2$. Let (q, F) be a condition in $\mathbb{Q}(\mathbb{P}_n)$ and suppose that for every σ which lies on (q, F) , we have a finite set \mathcal{X}_σ of conditions $p \leq q \restriction \sigma$. For every τ which lies on $(q \restriction n, F \restriction n)$, let \mathcal{X}_τ be the set of all conditions $p \restriction n$ with $p \in \mathcal{X}_\sigma$ for $\sigma \restriction n = \tau$. Using the inductive hypothesis for n , let (r, J) be the condition for the sets \mathcal{X}_τ . Fix ρ lying on (r, J) , and find τ_ρ which lies on $(q \restriction n, F \restriction n)$ such that ρ extends τ_ρ . Since (q, F) was determined, $r \restriction \rho$ decides the $F(n)$ -th level of $q(n)$. So let \dot{T}^ρ be a \mathbb{P}_n -name for the tree we get by replacing, for every node t on level $F(n)$ of $q(n)$, each $q(n)_t$ with the union of $p(n)$ for $p \in \mathcal{X}_\sigma$ with $\sigma = \tau_\rho \cup \{(n, t)\}$ and $r \restriction \rho \leq p \restriction n$. Strengthen $r \restriction \rho$ to r'_ρ deciding a level $m_\rho > F(n)$ such that for every node u on level m_ρ of \dot{T}^ρ above a node t on level $F(n)$, we can thin out \dot{T}_u^ρ to $\dot{P}^{\rho,u}$ such that $\dot{P}^{\rho,u} \leq p(n)$ for some $p \in \mathcal{X}_\sigma$ with $\sigma = \tau_\rho \cup \{(n, t)\}$ and $r \restriction \rho \leq p \restriction n$, and moreover for every such $p(n)$, there is some node u with $\dot{P}^{\rho,u} \leq p(n)$. We can assume, with some more thinning out, that all $m_\rho = m$ for a fixed m . Let (r', J') be the condition we get by applying the inductive hypothesis to (r, J) and $\mathcal{X}_\rho = \{r'_\rho\}$. Let \dot{T}'^ρ a \mathbb{P}_n -name for the tree we get by replacing, for every node u on level m of \dot{T}^ρ , each \dot{T}_u^ρ with $\dot{P}^{\rho,u}$. Let $\bar{F} = J' \cup \{(n, m)\}$. Let $X_{\bar{F}}$ consist of $\sigma : n + 1 \rightarrow {}^{<\omega}2$ such that $\sigma = \tau \cup \{(n, u)\}$ for some τ which lies on (r', J') with τ extending ρ lying on (r, J) and u on level m of \dot{T}'^ρ . Let φ be the $X_{\bar{F}}$ -assignment such that $\varphi(\sigma) = r' \restriction (\sigma \restriction n) \cup \{(n, \dot{T}'_{\sigma(n)}{}^\rho)\}$, where $\sigma \restriction n$ extends ρ lying on (r, J) . Let \bar{q} be the amalgamation of φ . Clearly the condition (\bar{q}, \bar{F}) has all the desired properties. \square

5. GROWING FINITE ITERATIONS OF PERFECT POSETS

In the construction of Jensen's poset \mathbb{P}^J , at nontrivial stages α , we used the ω -many perfect trees obtained from a partially generic filter for $\mathbb{Q}(\mathbb{P}_\alpha)^{<\omega}$ to grow the perfect poset \mathbb{P}_α to $\mathbb{P}_{\alpha+1}$. What we would like to do now is to find an appropriate generalization of this construction for growing a finite iteration \mathbb{P}_n of perfect posets using partially generic filters for the associated fusion poset $\mathbb{Q}(\mathbb{P}_n)$. More precisely, we would like the following.

Given a finite iteration $\mathbb{P}_n = \mathbb{Q}_0 * \dot{\mathbb{Q}}_1 * \cdots * \dot{\mathbb{Q}}_{n-1}$ of perfect posets, we would like to be able to extend it to a finite iteration $\mathbb{P}_n^* = \mathbb{Q}_0^* * \dot{\mathbb{Q}}_1^* * \cdots * \dot{\mathbb{Q}}_{n-1}^*$ of perfect posets, constructed in a generic extension of some suitable model M , with the following properties:

- (1) $\mathbb{Q}_0 \subseteq \mathbb{Q}_0^*$,

- (2) For all $1 \leq i < n$, $\mathbb{1}_{\mathbb{P}_i^*}$ forces that $\dot{\mathbb{Q}}_i$ is a perfect poset and $\dot{\mathbb{Q}}_i^*$ extends it.
- (3) $\mathbb{P}_n \subseteq \mathbb{P}_n^*$.
- (4) Every maximal antichain $\mathcal{A} \in M$ of \mathbb{P}_n remains maximal in \mathbb{P}_n^* .

The next theorem from [Abr84] holds the main idea for constructing \mathbb{P}_n^* . The set-up for the theorem is left intentionally vague with the details forthcoming in the next section.

Fix a suitable model M with $\mathbb{P}_n \in M$. We carry out the construction of \mathbb{P}_n^* in n -steps, at each step i , constructing a \mathbb{P}_i^* -name $\dot{\mathbb{Q}}_i^*$ for a perfect poset extending $\dot{\mathbb{Q}}_i$. We extend \mathbb{Q}_0 to \mathbb{Q}_0^* as before in a (carefully chosen) forcing extension of M by $\mathbb{Q}(\mathbb{Q}_0)^{<\omega}$. So suppose inductively that we already extended \mathbb{P}_i to \mathbb{P}_i^* satisfying requirements (1)-(4). Thus, in particular, every maximal antichain $\mathcal{A} \in M$ of \mathbb{P}_i remains maximal in \mathbb{P}_i^* , and so every V -generic filter H^* for \mathbb{P}_i^* restricts to an M -generic filter H for \mathbb{P}_i . From this it follows that $\dot{\mathbb{Q}}_i$ is forced to be a perfect poset by \mathbb{P}_i^* because $M[H]$ satisfies that $(\dot{\mathbb{Q}}_i)_H$ is a perfect poset, and this statement is clearly absolute between $M[H]$ and $V[H^*]$. Fix a carefully chosen M -generic filter $G \subseteq \mathbb{Q}(\mathbb{P}_{i+1})$. Let $\tau(G)$ be a \mathbb{P}_i^* -name for a subset of $\mathbb{Q}(\dot{\mathbb{Q}}_i)$ such that in any forcing extension $V[H^*]$ by \mathbb{P}_i^* $\tau(G)$ gets interpreted as the collection of all pairs $(p(i)_{H^*}, F(i))$ for $(p, F) \in G$. Provided that the poset \mathbb{P}_i^* contains a kind of master condition for G , we will be able to conclude that $\tau(G)$ is $M[H]$ -generic for $\mathbb{Q}((\dot{\mathbb{Q}}_i)_H)$, where H is the restriction of the V -generic filter $H^* \subseteq \mathbb{P}_i^*$.

Theorem 5.1. *Suppose $\bar{p} \in \mathbb{P}_i^*$ is such that for every $(p, F) \in G$, $\bar{p} \leq p \upharpoonright i$. Then*

$$\bar{p} \Vdash \text{“}\tau(G) \text{ is an } M[H]\text{-generic filter for } \mathbb{Q}(\dot{\mathbb{Q}}_i)\text{”},$$

where H is the restriction of the generic filter $H^* \subseteq \mathbb{P}_i^*$ to \mathbb{P}_i .

Proof. Suppose $H^* \subseteq \mathbb{P}_i^*$ is V -generic with $\bar{p} \in H^*$. Let H be the restriction of H^* to \mathbb{P}_i . Let

$$K = \tau(G)_{H^*} = \{(p(i)_{H^*}, F(i)) \mid p \in G\}$$

and $\mathbb{Q}_i = (\dot{\mathbb{Q}}_i)_H$. First, we argue that K is a filter on $\mathbb{Q}(\mathbb{Q}_i)$. Suppose for $(p, F) \in G$, $(p(i)_H, F(i)) \leq (T, b)$ in $\mathbb{Q}(\mathbb{Q}_i)$, so that $b \leq F(i)$ and $p(i)_H \cap b2 = T \cap b2$. It follows that there is a \mathbb{P}_i -name \dot{T} for T such that

$$\mathbb{1}_{\mathbb{P}_i} \Vdash \text{“}p(i) \leq \dot{T} \text{ and } p(i) \cap b2 = \dot{T} \cap b2\text{”}.$$

Let $p' = p \upharpoonright i \cup \{(i, \dot{T})\}$ and $F' = F \cup \{(i, b)\}$. Clearly $(p, F) \leq (p', F')$, which means that $(p', F') \in G$. It follows that $p'(i)_H = T$, and so $(T, b) \in K$.

Next, we fix (p, F) and (p', F') both in G and argue that $(p(i)_H, F(i))$ and $(p'(i)_H, F'(i))$ are compatible in K . There is $(q, J) \in G$ below both (p, F) and (p', F') . It follows that $J(i) \geq F(i), F'(i)$, $q \upharpoonright i \Vdash_{\mathbb{P}_i} q(i) \leq p(i), p'(i)$,

$$q \upharpoonright i \Vdash_{\mathbb{P}_i} \text{“}q(i) \cap F(i)2 = p(i) \cap F(i)2 \text{ and } q(i) \cap F'(i)2 = p'(i) \cap F'(i)2\text{”},$$

and $(q(i)_H, J(i)) \in K$. Since $\bar{p} \leq q \upharpoonright i$ and $\bar{p} \in H^*$, we have $q \upharpoonright i \in H^*$. Now observe that $q \upharpoonright i$ must force the statements mentioned above over \mathbb{P}_i^* as well because any V -generic filter for \mathbb{P}_i^* restricts to an M -generic filter for \mathbb{P}_i and the statements in question are absolute. Thus,

$$q(i)_H \leq p(i)_H, p'(i)_H$$

and

$$q(i)_H \cap F(i)2 = p(i)_H \cap F(i)2 \text{ and } q(i)_H \cap F'(i)2 = p'(i)_H \cap F'(i)2.$$

So $(q(i)_H, J(i)) \leq (p(i)_H, F(i))$, $(p'(i)_H, F'(i))$.

Finally, we have to see that K is $M[H]$ -generic. So suppose $D \in M[H]$ is dense open in $\mathbb{Q}(\mathbb{Q}_i)$. Let $\dot{D} \in M$ be a \mathbb{P}_i -name for D such that

$$\mathbb{1}_{\mathbb{P}_i} \Vdash_{\mathbb{P}_i} \text{“}\dot{D} \text{ is dense open in } \mathbb{Q}(\dot{\mathbb{Q}}_i)\text{”}.$$

In M , define

$$E = \{(p, F) \in \mathbb{Q}(\mathbb{P}_{i+1}) \mid p \upharpoonright i \Vdash (p(i), F(i)) \in \dot{D}\}.$$

We claim that E is dense open in $\mathbb{Q}(\mathbb{P}_{i+1})$. It is easy to see that E is open, so let's argue that it is dense. Fix some $(q, J) \in \mathbb{Q}(\mathbb{P}_{i+1})$ and assume without loss that (q, J) is determined. There must be some pair (\dot{T}, \dot{k}) such that \dot{T} is a \mathbb{P}_i -name for an element of $\dot{\mathbb{Q}}_i$, \dot{k} is a \mathbb{P}_i -name for a natural number, and

$$q \upharpoonright i \Vdash \text{“}(\dot{T}, \dot{k}) \in \dot{D} \text{ and } (\dot{T}, \dot{k}) \leq (q(i), J(i))\text{”}.$$

The set of conditions which decide the value of \dot{k} is dense open below $q \upharpoonright i$ in \mathbb{P}_i . So, by Proposition 4.4, there a determined condition $(p', F') \leq (q \upharpoonright i, J \upharpoonright i)$ in $\mathbb{Q}(\mathbb{P}_i)$ such that for every σ which lies on it, $p' \upharpoonright \sigma$ decides that $\dot{k} = k(\sigma)$. Let $k \in \omega$ be above all the $k(\sigma)$. Then

$$p' \Vdash \text{“}(\dot{T}, k) \leq (\dot{T}, \dot{k}) \leq (q(i), J(i)) \text{ and } (\dot{T}, k) \in \dot{D}\text{”}.$$

Let $p = p' \cup \{(i, \dot{T})\}$ and $F = F' \cup \{(i, k)\}$. Clearly $(p, F) \in E$. Let's argue that $(p, F) \leq (q, J)$. By construction $p \leq q$ and each $k(\sigma) \geq J(i)$, so $k \geq J(i)$. Finally, $p' = p \upharpoonright i \Vdash (\dot{T}, k) \leq (q(i), J(i))$ and so $p \upharpoonright i \Vdash \dot{T} \cap J(i)2 = q(i) \cap J(i)2$.

Fix some $(p, F) \in E \cap G$. Since $\bar{p} \leq p \upharpoonright i$, it follows that $p \upharpoonright i \in H^*$. Thus, $(p(i)_H, F(i)) \in D \cap K$, completing the argument that K is an $M[H]$ -generic filter for $\mathbb{Q}(\mathbb{Q}_i)$. \square

Next, we are going to obtain a stronger version of Theorem 5.1 that tells us how to get an $M[H]$ -generic filter for $\mathbb{Q}(\mathbb{Q}_i)^{<\omega}$, which is really what we need to extend $\dot{\mathbb{Q}}_i$ to $\dot{\mathbb{Q}}_i^*$. For this, we will need to enlarge our fusion poset $\mathbb{Q}(\mathbb{P}_{i+1})$. Let $\bar{\mathbb{Q}}_i(\mathbb{P}_{i+1})$ be the following modification of $\mathbb{Q}(\mathbb{P}_{i+1})$. Conditions in $\bar{\mathbb{Q}}_i(\mathbb{P}_{i+1})$ are pairs (p, F) such that $(p \upharpoonright i, F \upharpoonright i) \in \mathbb{Q}(\mathbb{P}_i)$, $p(i)$ is some finite tuple $\langle \dot{T}_0, \dot{T}_1, \dots, \dot{T}_{k-1} \rangle$ with $p \upharpoonright i \Vdash \dot{T}_j \in \dot{\mathbb{Q}}_i$ for all $j < k$, and $F(i) = f : k \rightarrow \omega$. The ordering is $(p', F') \leq (p, F)$ whenever

- (1) $(p' \upharpoonright i, F' \upharpoonright i) \leq (p \upharpoonright i, F \upharpoonright i)$, and
for $j < k$,
- (2) $F'(i)(j) \geq F(i)(j)$,
- (3) $p' \upharpoonright i \Vdash p'(i)(j) \cap F'(i)(j)2 = p(i)(j) \cap F(i)(j)2$.

The point is that if $H^* \subseteq \mathbb{P}_i^*$ is V -generic, then

$$\langle ((\dot{T}_0)_{H^*}, f(0)), \dots, ((\dot{T}_{k-1})_{H^*}, f(k-1)) \rangle$$

is a condition in $\mathbb{Q}((\dot{\mathbb{Q}}_i)_H)^{<\omega}$. Suppose now that $G \subseteq \bar{\mathbb{Q}}_i(\mathbb{P}_{i+1})$ is M -generic. Let $\tau(G)$ be a \mathbb{P}_i^* -name for a subset of $\mathbb{Q}(\mathbb{Q}_i)^{<\omega}$ such that in any forcing extension $V[H^*]$ by \mathbb{P}_i^* , $\tau(G)$ gets interpreted as the collection of all

$$\langle ((\dot{T}_0)_{H^*}, f(0)), \dots, ((\dot{T}_{k-1})_{H^*}, f(k-1)) \rangle$$

for $(p, F) \in G$. If it so happens that \mathbb{P}_i^* has a master condition \bar{p} for G , then $\tau(G)$ will be $M[H]$ -generic for $\mathbb{Q}((\mathbb{Q}_i)_H)^{<\omega}$, where H is the restriction of the V -generic filter $H^* \subseteq \mathbb{P}_i^*$.

Theorem 5.2. *Suppose $\bar{p} \in \mathbb{P}_i^*$ is such that for every $(p, F) \in G$, $\bar{p} \leq p \upharpoonright i$. Then*

$$\bar{p} \Vdash \text{“}\tau(G) \text{ is an } M[H]\text{-generic filter for } \mathbb{Q}(\dot{\mathbb{Q}}_i)^{<\omega}\text{”},$$

where H^* is the restriction of the generic filter to \mathbb{P}_i .

The proof is essentially the same as of Theorem 5.1. Using Theorem 5.2, we can let $\dot{\mathbb{Q}}_i^*$ be the canonical \mathbb{P}_i^* -name for the extension of $\dot{\mathbb{Q}}_i$ formed in $M[H]$ from $\tau(G)$. In the next section, we will show how to obtain the required M -generic filters G so that the inductive assumptions hold for \mathbb{P}_i^* .

6. TREE ITERATIONS OF PERFECT POSETS

Let's define that an ω -iteration of perfect posets is a sequence

$$\vec{P} = \langle \mathbb{P}_n \mid n < \omega \rangle,$$

where $\mathbb{P}_0 = \{\emptyset\}$ is a trivial poset and each \mathbb{P}_n , for $n \geq 1$, is a finite iteration of perfect posets with the coherence requirement that for $0 < m < n$, $\mathbb{P}_n \upharpoonright m = \mathbb{P}_m$. We included \mathbb{P}_0 in the sequence to make the subsequent definitions more uniform. For this reason, we will also make the ad hoc definition that $\mathbb{Q}(\mathbb{P}_0) = \{\emptyset\}$. Note that an ω -iteration of perfect posets is not itself an iteration, rather it is a coherent sequence of finite iterations.

A tree iteration is a non-linear iteration along some tree. Given a tree of height ω , the tree iteration of perfect posets will use an ω -iteration $\vec{P} = \langle \mathbb{P}_n \mid n < \omega \rangle$ of perfect posets with conditions assigned to level n nodes of the tree coming from \mathbb{P}_n . Conditions will be assigned to the nodes coherently so that if a node s on level n extends a node t on level m , then the condition p on node s will be such that $p \upharpoonright m$ is on node t .

Definition 6.1. Let $\vec{P} = \langle \mathbb{P}_n \mid n < \omega \rangle$ be an ω -iteration of perfect posets and let \mathcal{T} be a tree of height ω . A \mathcal{T} -iteration of perfect posets is the following partial order $\mathbb{P}(\vec{P}, \mathcal{T})$. Conditions in $\mathbb{P}(\vec{P}, \mathcal{T})$ are functions f_X with domain some finite subtree X of \mathcal{T} such that:

- (1) For every node s on level n of X , $f_X(s) \in \mathbb{P}_n$.
- (2) Whenever $s \leq t$ are two nodes in X , then $f_X(t) \upharpoonright \text{lev}(s) = f_X(s)$.

The ordering is $f_Y \leq f_X$ whenever Y extends X and for every node $s \in X$, $f_Y(s) \leq f_X(s)$.

The analogue of the fusion posets $\mathbb{Q}(\mathbb{P})$ and $\mathbb{Q}(\mathbb{P}_n)$ for $\mathbb{P}(\vec{P}, \mathcal{T})$ will be the fusion poset $\mathbb{Q}(\vec{P}, \mathcal{T})$ whose conditions are functions f_X with domain some finite subtree X of \mathcal{T} such that:

- (1) For every node s on level n of X , $f_X(s) \in \mathbb{Q}(\mathbb{P}_n)$.
- (2) Whenever $s \leq t$ are two nodes in X , with $f_X(s) = (p_s, F_s)$ and $f_X(t) = (p_t, F_t)$, then $p_t \upharpoonright \text{lev}(s) = p_s$ and $F_t \upharpoonright \text{lev}(s) = F_s$.

The ordering is $f_Y \leq f_X$ whenever Y extends X and for every node $s \in X$, $f_Y(s) \leq f_X(s)$.

Proposition 6.2.

- (1) Suppose $f_X \in \mathbb{P}(\vec{P}, \mathcal{T})$, $f_X(s) = p$, and $q \leq p$. Then there is a condition $g_X \leq f_X$ in $\mathbb{P}(\vec{P}, \mathcal{T})$ with $g_X(s) = q$.
- (2) Suppose $f_X \in \mathbb{Q}(\vec{P}, \mathcal{T})$, $f_X(s) = (p, F)$, and $(q, F') \leq (p, F)$. Then there is $g_X \leq f_X$ such that $g_X(s) = (q, F')$.

Proof. We will only prove (1) because the proof of (2) is essentially the same. Define g_X as follows. Fix a node $t \in X$. If $t \leq s$, then let $g_X(t) = q \upharpoonright \text{lev}(t)$. If $s \leq t$, then let $g_X(t)$ be q concatenated with the tail of $f_X(t)$. Otherwise, let t' be the largest node that is compatible with both t and s . Let $g_X(t)$ be $q \upharpoonright \text{lev}(t')$ concatenated with the tail of $f_X(t)$. \square

Let us call a condition $f_X \in \mathbb{Q}(\vec{P}, \mathcal{T})$ *determined* if every $f_X(s)$ is determined. Note that to check whether a condition f_X is determined, it suffices to check that the conditions on the terminal nodes are determined.

Proposition 6.3. *The set of all determined conditions is dense in $\mathbb{Q}(\vec{P}, \mathcal{T})$.*

Proof. Fix $f_X \in \mathbb{Q}(\vec{P}, \mathcal{T})$. Let $\langle s_i \mid i \leq m \rangle$ be an enumeration of the terminal nodes of X . Using the construction in the proof of Proposition 6.2, strengthen f_X to f_X^0 such that $f_X^0(s_0)$ is determined. Inductively, given f_X^i , we let f_X^{i+1} be the strengthening of f_X^i constructed as in the proof of Proposition 6.2 such that $f_X^{i+1}(s_{i+1})$ is determined. Let's argue that f_X^m is determined. We can assume inductively that the conditions $f_X^{m-1}(s_i)$ for $i < m$ are determined. By construction $f_X^m(s_m)$ is determined. So fix $i < m$. Let s be the node where branch of s_i and the branch of s_m split. By construction, $f_X^m(s_i)$ is $f_X^m(s_m) \upharpoonright \text{lev}(s)$ concatenated with the tail of $f_X^{m-1}(s_i)$. Clearly $f_X^m(s_m) \upharpoonright \text{lev}(s)$ is determined and $f_X^m(s_m) \upharpoonright \text{lev}(s) \leq f_X^{m-1}(s_i) \upharpoonright \text{lev}(s)$. Now using the definition of what it means to be determined, it is easy to see that the whole condition is determined. \square

We will initially consider tree iterations along the countable tree ${}^{<\omega}\omega$, and then later extend our results to tree iterations along the uncountable tree ${}^{<\omega}\omega_1$.

It is easy to see that the poset $\mathbb{Q}(\mathbb{Q}_0)^{<\omega}$ completely embeds into $\mathbb{Q}(\vec{P}, {}^{<\omega}\omega)$ via the map sending a condition to the corresponding tree of height ≤ 2 .

Proposition 6.4. *The poset $\mathbb{Q}(\mathbb{Q}_0)^{<\omega}$ completely embeds into $\mathbb{Q}(\vec{P}, {}^{<\omega}\omega)$ via the following map φ :*

- (1) $\varphi(\mathbb{1}_{\mathbb{Q}(\mathbb{Q}_0)}) = f_X$, where X consists of the root node s and $f_X(s) = \emptyset$.
For p with some non-trivial $p(i)$,
- (2) $\varphi(p) = f_X$, where X consists of the root node together with nodes $\langle i \rangle$ for non-trivial $p(i)$, such that $f_X(\langle i \rangle) = p(i)$.

More generally, for each node s on level n of ${}^{<\omega}\omega$, $\bar{\mathbb{Q}}(\mathbb{P}_{n+1})$ completely embeds into $\mathbb{Q}(\vec{P}, {}^{<\omega}\omega)$ via the map sending a condition to the corresponding tree of height $\leq n+2$ whose stem stretches up to s .

Proposition 6.5. *Fix a node s on level n in ${}^{<\omega}\omega$. The poset $\bar{\mathbb{Q}}(\mathbb{P}_{n+1})$ completely embeds into $\mathbb{Q}(\vec{P}, {}^{<\omega}\omega)$ via the following map φ_s :*

- (1) $\varphi_s(\mathbb{1}_{\bar{\mathbb{Q}}(\mathbb{P}_{n+1})}) = f_X$, where X is the branch ending in s , such that $f_X(s) = \mathbb{1}_{\mathbb{Q}(\mathbb{P}_n)}$.
For (p, F) with $\text{dom}(p(n)) = k$,
- (2) $\varphi_s((p, F)) = f_X$, where X consists of the branch ending in s together with nodes $\langle s \hat{\ } i \rangle$ for $i < k$, such that $f_X(s) = (p \upharpoonright n, F \upharpoonright n)$ and
 $f_X(s \hat{\ } i) = (p \upharpoonright n \hat{\ } p(n)(i), F \upharpoonright n \hat{\ } F(n)(i))$.

Suppose $G \subseteq \mathbb{Q}(\vec{P}, {}^{<\omega}\omega)$ is V -generic and fix some node s on level n of ${}^{<\omega}\omega$. We will use the notation G_s for the V -generic filter for $\bar{\mathbb{Q}}(\mathbb{P}_{n+1})$ added by G via the

embedding φ_s and we will use the notation G_\emptyset for the V -generic filter for $\mathbb{Q}(\mathbb{Q}_0)^{<\omega}$ added by G via the embedding φ .

Suppose now that $\vec{P} = \langle \mathbb{P}_n \mid n < \omega \rangle$ is an ω -iteration of perfect posets that is an element of a suitable model M . Let $G \subseteq \mathbb{Q}(\vec{P}, <^\omega \omega)$ be M -generic. We will argue that in $M[G]$, we can grow each iteration \mathbb{P}_n to an iteration \mathbb{P}_n^* satisfying requirements (1)-(4) from Section 5:

- (1) $\mathbb{Q}_0 \subseteq \mathbb{Q}_0^*$,
- (2) For all $1 \leq i < n$, $\mathbb{1}_{\mathbb{P}_i^*}$ forces that $\dot{\mathbb{Q}}_i$ is a perfect poset and $\dot{\mathbb{Q}}_i^*$ extends it.
- (3) $\mathbb{P}_n \subseteq \mathbb{P}_n^*$.
- (4) Every maximal antichain $\mathcal{A} \in M$ of \mathbb{P}_n remains maximal in \mathbb{P}_n^* .

It is straightforward to extend \mathbb{Q}_0 to \mathbb{Q}_0^* . By Proposition 6.4, G adds an M -generic filter G_\emptyset for $\mathbb{Q}(\mathbb{Q}_0)^{<\omega}$. Let \mathcal{T}_i^0 for $i < \omega$ be the generic perfect trees added by G_\emptyset and construct \mathbb{Q}_0^* as before. Recall that $\langle \mathcal{T}_i^0 \mid i < \omega \rangle$ is a maximal antichain in \mathbb{Q}_0^* and every maximal antichain $\mathcal{A} \in M$ of \mathbb{Q}_0 remains maximal in \mathbb{Q}_0^* .

Now let's show how to extend $\dot{\mathbb{Q}}_1$ to a \mathbb{P}_1^* -name $\dot{\mathbb{Q}}_1^*$ for a perfect poset. By Proposition 6.5, for each node s on level 1 of the tree $<^\omega \omega$, G adds an M -generic filter G_s for $\mathbb{Q}(\mathbb{P}_2)$. Observe that each $\mathcal{T}_i^0 \leq p \upharpoonright 1$ for all p with $(p, F) \in G_{\langle i \rangle}$. Thus, by Theorem 5.2, whenever $H^* \subseteq \mathbb{P}_1^*$ is a V -generic filter containing \mathcal{T}_i^0 , then the interpretation $\tau(G_{\langle i \rangle})_{H^*}$ is an $M[H]$ -generic filter for $\mathbb{Q}((\dot{\mathbb{Q}}_1)_H)^{<\omega}$, where H is the restriction of H^* to \mathbb{P}_1 . Let τ be a mixed \mathbb{P}_1^* -name that is interpreted as $\tau(G_{\langle i \rangle})_{H^*}$, whenever $\mathcal{T}_i^0 \in H^*$. Since the conditions \mathcal{T}_i^0 form a maximal antichain in \mathbb{P}_1^* ,

$$\mathbb{1}_{\mathbb{P}_1^*} \Vdash \text{“}\tau \text{ is an } M[H]\text{-generic filter for } \mathbb{Q}(\dot{\mathbb{Q}}_1)^{<\omega}\text{”}.$$

So let $\dot{\mathbb{Q}}_1^*$ be a \mathbb{P}_1^* -name for the perfect poset constructed as usual from $\dot{\mathbb{Q}}_1$ and τ .

For each $j < \omega$, we can choose a \mathbb{P}_1^* -name $\dot{\mathcal{T}}_j^1$ for the perfect tree on coordinate j of τ . The pairs $(\mathcal{T}_i^0, \dot{\mathcal{T}}_j^1)$ form a maximal antichain in \mathbb{P}_2^* because the collection of \mathcal{T}_i^0 is a maximal antichain and each \mathcal{T}_i^0 forces that the trees $\dot{\mathcal{T}}_j^1$ form a maximal antichain in $\dot{\mathbb{Q}}_1^*$. Also, clearly for every $(p, F) \in G_{\langle i, j \rangle}$, we have $(\mathcal{T}_i^0, \dot{\mathcal{T}}_j^1) \leq p \upharpoonright 2$. This will allow us to apply Theorem 5.2 to grow $\dot{\mathbb{Q}}_2$ to $\dot{\mathbb{Q}}_2^*$. Finally, let's argue that \mathbb{P}_2 is actually a subset of \mathbb{P}_2^* . Suppose p is a condition in \mathbb{P}_2 . Then $p(0) \in \mathbb{P}_1$ and hence $p(0) \in \mathbb{P}_1^*$ as well. Also, clearly $p(1)$ is a \mathbb{P}_1^* -name. So we need to argue that $p(0) \Vdash_{\mathbb{P}_1^*} p(1) \in \dot{\mathbb{Q}}_1^*$. Fix a V -generic filter $H^* \subseteq \mathbb{P}_1^*$ with $p(0) \in H^*$ and consider $M[H]$ where H is the restriction of H^* to \mathbb{P}_1 . Since $p(0) \Vdash_{\mathbb{P}_1} p(1) \in \dot{\mathbb{Q}}_1$, in $M[H]$, we have $p(1)_H \in (\dot{\mathbb{Q}}_1)_H \subseteq (\dot{\mathbb{Q}}_1^*)_{H^*}$. It remains to show (4), that every maximal antichain of \mathbb{P}_2 from M remains maximal in \mathbb{P}_2^* . We will provide an inductive proof of this later in Lemma 6.8.

For now to finish the construction, we assume that properties (1)-(4) hold for \mathbb{P}_n^* . We will additionally assume that:

- (1) For each node s on level n of $<^\omega \omega$ there is a condition

$$(\mathcal{T}_{s(0)}^0, \dot{\mathcal{T}}_{s(1)}^1, \dots, \dot{\mathcal{T}}_{s(n-1)}^{n-1}) \in \mathbb{P}_n^*$$

which is below all $p \upharpoonright n$ with $(p, F) \in G_s$.

- (2) The collection of all such conditions $(\mathcal{T}_{s(0)}^0, \dot{\mathcal{T}}_{s(1)}^1, \dots, \dot{\mathcal{T}}_{s(n-1)}^{n-1})$ is a maximal antichain in \mathbb{P}_n^* .

With this set-up, we extend $\dot{\mathbb{Q}}_n$ to $\dot{\mathbb{Q}}_n^*$ identically to the case $n = 1$ above, using Theorem 5.2. It is also easy to see inductively that \mathbb{P}_n is a subset of \mathbb{P}_n^* .

To get some intuition for the construction, let's fix an L -generic filter $H^* \subseteq \mathbb{P}_1^*$ and see what a condition $\mathcal{T}_j^1 = (\dot{\mathcal{T}}_j^1)_{H^*}$ looks like. Since $\langle \mathcal{T}_i^0 \mid i < \omega \rangle$ is a maximal antichain of \mathbb{P}_1^* , there is a unique i such that $\mathcal{T}_i^0 \in H^*$. Let

$$K = \{(p(1)_{H^*}, F(1)) \mid f_X \in G \text{ and } f_X(\langle i, j \rangle) = (p, F)\}.$$

Then \mathcal{T}_j^1 is the union of $T \cap {}^{n_2}$ for $(T, n) \in K$.

To give the promised argument that every maximal antichain of \mathbb{P}_n from M remains maximal in \mathbb{P}_n^* , we first need to define the analogue of $\mathbb{U} \subseteq \mathbb{P}^*$ from Section 2 for \mathbb{P}_n^* .

Let \mathbb{U}_n be the subset of \mathbb{P}_n^* consisting of conditions $p = (p_0, \dot{p}_1, \dots, \dot{p}_{n-1})$ such that $p_0 = \mathcal{T}_{j_0}^0 \wedge S_0$ for some $j_0 < \omega$ and $S_0 \in \mathbb{Q}_0$, and for $i < n$, $p \upharpoonright i \Vdash \dot{p}_i = \dot{\mathcal{T}}_{j_i}^i \wedge \dot{S}_i$ for some $j_i < \omega$ and a \mathbb{P}_i -name \dot{S}_i such that $\mathbb{1}_{\mathbb{P}_i} \Vdash \dot{S}_i \in \dot{\mathbb{Q}}_i$.

Proposition 6.6. \mathbb{U}_n is dense in \mathbb{P}_n^* .

Proof. We argue by induction on n . By Proposition 2.4, \mathbb{U}_1 is dense in \mathbb{P}_1^* and every maximal antichain of \mathbb{P}_1 from M remains maximal in \mathbb{P}_1^* . So let's suppose inductively that \mathbb{U}_n is dense in \mathbb{P}_n^* and every maximal antichain of \mathbb{P}_n from M remains maximal in \mathbb{P}_n^* . This argument is meant to take place simultaneously with the inductive proof of Lemma 6.8, where we use the density of \mathbb{U}_n to argue that maximal antichains stay maximal.

Fix a condition $(q'_0, \dot{q}'_1, \dots, \dot{q}'_n)$ in \mathbb{P}_{n+1}^* . In particular, we have $(q'_0, \dot{q}'_1, \dots, \dot{q}'_{n-1}) \Vdash \dot{q}'_n \in \mathbb{Q}_n^*$. Consider any forcing extension $V[H^*]$ by \mathbb{P}_n^* with $(q'_0, \dot{q}'_1, \dots, \dot{q}'_{n-1}) \in H^*$. In $V[H^*]$, there is $j_n < \omega$ and $S \in (\dot{\mathbb{Q}}_n)_{H^*}$ such that $(\dot{\mathcal{T}}_{j_n}^n)_{H^*} \wedge S \leq (q'_n)_{H^*}$. So $S \in M[H]$, where H is the restriction of H^* to \mathbb{P}_n . Let \dot{S} be a \mathbb{P}_n -name in M such that $(\dot{S})_H = S$ and let $p \in H$ force that $\dot{S} \in \dot{\mathbb{Q}}_n$. Let $\mathcal{A} \in M$ be any maximal antichain of \mathbb{P}_n extending $\{p\}$ and let \dot{S}_n be the mixed name such that $p \Vdash \dot{S} = \dot{S}_n$ and every other $q \in \mathcal{A}$ forces that $\dot{S}_n = {}^{<\omega}2$. So $\mathbb{1}_{\mathbb{P}_n} \Vdash \dot{S}_n \in \dot{\mathbb{Q}}_n$. Now let $(q_0, \dot{q}_1, \dots, \dot{q}_{n-1}) \leq (q'_0, \dot{q}'_1, \dots, \dot{q}'_{n-1})$ be a condition in H^* forcing that $\dot{\mathcal{T}}_{j_n}^n \wedge \dot{S}_n \leq q'_n$. We have just argued that there is a condition $(q_0, \dot{q}_1, \dots, \dot{q}_{n-1}) \leq (q'_0, \dot{q}'_1, \dots, \dot{q}'_{n-1})$ and a \mathbb{P}_n -name \dot{S}_n such that $\mathbb{1}_{\mathbb{P}_n} \Vdash \dot{S}_n \in \dot{\mathbb{Q}}_n$ and

$$(q_0, \dot{q}_1, \dots, \dot{q}_{n-1}) \Vdash \dot{\mathcal{T}}_{j_n}^n \wedge \dot{S}_n \leq q'_n$$

for some $j_n < \omega$.

Now, by the inductive assumption, there is a condition

$$(p_0, \dot{p}_1, \dots, \dot{p}_{n-1}) \leq (q_0, \dot{q}_1, \dots, \dot{q}_{n-1})$$

in \mathbb{U}_n . So clearly the condition $(p_0, \dot{p}_1, \dots, \dot{p}_{n-1}, \dot{\mathcal{T}}_{j_n}^n \wedge \dot{S}_n) \in \mathbb{U}_{n+1}$. \square

We will abuse notation by writing conditions in \mathbb{U}_n in the form

$$(\mathcal{T}_{j_0}^0 \wedge S_0, \dot{\mathcal{T}}_{j_1}^1 \wedge \dot{S}_1, \dots, \dot{\mathcal{T}}_{j_{n-1}}^{n-1} \wedge \dot{S}_{n-1}).$$

The next lemma is a generalization of Proposition 2.3.

Lemma 6.7. Suppose $f_X \in \mathbb{Q}(\vec{P}, {}^{<\omega}\omega)$ forces that

$$(\mathcal{T}_{j_0}^0 \wedge S_0, \dot{\mathcal{T}}_{j_1}^1 \wedge \dot{S}_1, \dots, \dot{\mathcal{T}}_{j_{n-1}}^{n-1} \wedge \dot{S}_{n-1}) \in \mathbb{U}_n.$$

Then there is a condition $g_Y \leq f_X$ such that $g_Y(\langle j_0, j_1, \dots, j_{n-1} \rangle) = (q, F)$ is determined and $\tau : n \rightarrow {}^{<\omega}2$ lies on (q, F) such that

$$q(0)_{\tau(0)} \leq S_0, \text{ and for all } 1 \leq i < n, (q \upharpoonright i) \mid (\tau \upharpoonright i) \Vdash q(i)_{\tau(i)} \leq \dot{S}_i.$$

Proof. The case $n = 1$ follows by Proposition 2.3. So suppose $n = 2$. Recall that every maximal antichain of \mathbb{P}_1 from M remains maximal in \mathbb{P}_1^* . By strengthening f_X , we can assume that $\langle j_0, j_1 \rangle \in X$. Let $f_X(\langle j_0, j_1 \rangle) = (\bar{q}, \bar{F})$. By strengthening further, using the case $n = 1$, we can assume that there is a node s_0 on level $\bar{F}(0)$ such that $\bar{q}(0)_{s_0} \leq S_0$. By strengthening some more, we can assume that (\bar{q}, \bar{F}) is determined.

Let's argue that $\bar{q}(0)_{s_0}$ forces that for some node s_1 on level $\bar{F}(1)$ of $\bar{q}(1)$, $\bar{q}(1)_{s_1} \wedge \dot{S}_1 \neq \emptyset$. If this is not the case, then there is some $T \leq \bar{q}(0)_{s_0}$ which forces that there is no such node. Let T' be the tree we get by replacing $\bar{q}(0)_{s_0}$ with T in $\bar{q}(0)$. Let $\bar{q}' = (T', \bar{q}(1))$ and let f'_X be a condition strengthening f_X so that $f'_X(\langle j_0, j_1 \rangle) = (\bar{q}', \bar{F})$. Let G be an M -generic filter for $\mathbb{Q}(\bar{P}, <^\omega \omega)$ containing f'_X and consider $M[G]$. The tree $R = \mathcal{T}_{j_0}^0 \wedge S_0 \wedge \bar{q}'(0)_{s_0} (= T)$ is a condition in \mathbb{P}_1^* , and so we can fix some V -generic filter $H^* \subseteq \mathbb{P}_1^*$ containing R . In particular, H^* contains $\mathcal{T}_{j_0}^0 \wedge S_0$ and T . Thus, in $M[H]$ (where H is the restriction of H^* to \mathbb{P}_1), $\bar{q}(1)_H$ does not have any subtree on level $F(1)$ which has a common subtree with $(\dot{S}_1)_H$. But then $(\dot{\mathcal{T}}_{j_1}^1)_{H^*} \wedge (\dot{S}_1)_H = \emptyset$ contradicting our assumption that f_X forced $(\mathcal{T}_{j_0}^0 \wedge S_0, \dot{\mathcal{T}}_{j_1}^1 \wedge \dot{S}_1) \in \mathbb{U}_2$.

Let's strengthen $\bar{q}(0)_{s_0}$ to some T_0 , so that there is a node s_1 on level $\bar{F}(1)$ such that $T_0 \Vdash \bar{q}(1)_{s_1} \wedge \dot{S}_1 \neq \emptyset$. Let \dot{T}_1 be a \mathbb{P}_1 -name such that

$$T_0 \Vdash \dot{T}_1 = \bar{q}(1)_{s_1} \wedge \dot{S}_1.$$

Since (\bar{q}, \bar{F}) is determined, it follows that $\bar{q}(0)_{s_0} \Vdash s_1 \in \bar{q}(1)$. Thus, $\tau = \{(0, s_0), (1, s_1)\}$ lies on (\bar{q}, \bar{F}) and $(T_0, \dot{T}_1) \leq \bar{q} \restriction \tau$. By Proposition 4.8, there is a determined condition $(q, F) \leq (\bar{q}, \bar{F})$ with ρ lying on (\bar{q}, \bar{F}) and extending τ such that $q \restriction \rho \leq (T_0, \dot{T}_1)$. Strengthen f_X to a condition g_X with $g_X(\langle j_0, j_1 \rangle) = (q, F)$ using Proposition 6.2 (2).

Suppose inductively that the statement holds for n and every maximal antichain of \mathbb{P}_n from M remains maximal in \mathbb{P}_n^* . Let's argue that the statement holds for $n+1$. This will basically be a generalization of the argument passing from $n = 1$ to $n = 2$. By strengthening f_X , we can assume that $\langle j_0, \dots, j_n \rangle \in X$ and that f_X is determined. Let $f_X(\langle j_0, \dots, j_n \rangle) = (\bar{q}, \bar{F})$. By using our inductive assumption for n and strengthening further, we can assume that $\tau : n \rightarrow <^\omega 2$ lies on $(\bar{q} \restriction n, \bar{F} \restriction n)$ such that

$$q(0)_{\tau(0)} \leq S_0, \text{ and for all } 1 \leq i < n, (q \restriction i) \restriction (\tau \restriction i) \Vdash q(i)_{\tau(i)} \leq \dot{S}_i.$$

Let's argue that $(\bar{q} \restriction n) \restriction \tau$ forces that for some node s_n on level $\bar{F}(n)$ of $\bar{q}(n)$, $\bar{q}(n)_{s_n} \wedge \dot{S}_n \neq \emptyset$. If this is not the case, then there is a condition $p \leq (\bar{q} \restriction n) \restriction \tau$ which forces that there is no such node. By Proposition 4.8, there is a determined condition $(p', F') \leq (\bar{q} \restriction n, \bar{F} \restriction n)$ with ρ lying on (p', F') and extending τ such that $p' \restriction \rho \leq p$. Let $\bar{q}' = p' \cup \{(n, \bar{q}(n))\}$ and $\bar{F}' = F' \cup \{(n, \bar{F}(n))\}$. Let f'_X be the strengthening of f_X in which $f'_X(\langle j_0, \dots, j_n \rangle) = (\bar{q}', \bar{F}')$. Let G be an M -generic filter for $\mathbb{Q}(\bar{P}, <^\omega \omega)$ containing f'_X and consider $M[G]$. The tree

$$R = (\mathcal{T}_{j_0}^0 \wedge S_0 \wedge \bar{q}'(0)_{\tau(0)}, \dots, \dot{\mathcal{T}}_{j_{n-1}}^{n-1} \wedge \dot{S}_{n-1} \wedge \bar{q}'(n-1)_{\tau(n-1)})$$

is a condition in \mathbb{P}_n^* (where $(\bar{q}'(0)_{\tau(0)}, \dots, \bar{q}'(n-1)_{\tau(n-1)}) \leq p$ by construction), and so we can fix a V -generic filter H^* containing R , and argue as in the case $n = 2$ towards a contradiction.

Thus, there is a condition

$$(T_0, \dot{T}_1, \dots, T_{n-1}) \leq (\bar{q} \upharpoonright n) \upharpoonright \tau,$$

a \mathbb{P}_n -name \dot{T}_n for a perfect tree, and a node s_n on level $\bar{F}(n)$ such that

$$(T_0, \dot{T}_1, \dots, T_{n-1}) \Vdash \dot{T}_n = \bar{q}(n)_{s_n} \wedge \dot{S}_n.$$

Let $\sigma = \tau \cup \{(n, s_n)\}$. So $(T_0, \dot{T}_1, \dots, \dot{T}_{n-1}, \dot{T}_n) \leq \bar{q} \upharpoonright \sigma$, and it remains to construct the required condition using Proposition 4.8. \square

We are now ready to prove that every maximal antichain of \mathbb{P}_n from M remains maximal in \mathbb{P}_n^* .

Lemma 6.8. *Every maximal antichain of \mathbb{P}_n from M remains maximal in \mathbb{P}_n^* .*

Proof. The statement is true for $n = 1$. So we can assume inductively that \mathbb{U}_n is dense in \mathbb{P}_n^* . Let $\mathcal{A} \in M$ be a maximal antichain in \mathbb{P}_n . It suffices to show that every condition in \mathbb{U}_n is compatible with an element of \mathcal{A} . So let f_X be some condition which forces that

$$(\mathcal{T}_{j_0}^0 \wedge S_0, \dot{T}_{j_1}^1 \wedge \dot{S}_1, \dots, \dot{T}_{j_{n-1}}^{n-1} \wedge \dot{S}_{n-1}) \in \mathbb{U}_n.$$

By Proposition 6.7, we can strengthen f_X to a condition g_Y such that $g_Y(\langle j_0, \dots, j_{n-1} \rangle) = (q, F)$ is determined and $\tau : n \rightarrow {}^{<\omega}2$ lies on (q, F) such that $q(0)_{\tau(0)} \leq S_0$, and for all $1 \leq i < n$, $(q \upharpoonright i) \upharpoonright (\tau \upharpoonright i) \Vdash q(i)_{\tau(i)} \leq \dot{S}_i$. Since \mathcal{A} is maximal in \mathbb{P}_n , $q \upharpoonright \tau$ is compatible with some $p \in \mathcal{A}$. So let $q' \leq q \upharpoonright \tau, p$. By Proposition 4.8, there is a determined condition $(\bar{q}, \bar{F}) \leq (q, F)$ with ρ lying on (\bar{q}, \bar{F}) and extending τ such that $\bar{q} \upharpoonright \rho \leq q'$. Strengthen g_Y to \bar{g}_Y with (q, F) strengthened to (\bar{q}, \bar{F}) as in Proposition 6.2 (2). The condition \bar{g}_Y forces that $(\mathcal{T}_{j_0}^0 \wedge S_0, \dot{T}_{j_1}^1 \wedge \dot{S}_1, \dots, \dot{T}_{j_{n-1}}^{n-1} \wedge \dot{S}_{n-1})$ is compatible with $p \in \mathcal{A}$. \square

Let $\vec{P}^* = \langle \mathbb{P}_n^* \mid n < \omega \rangle$ be the ω -iteration made up of the extended iterations \mathbb{P}_n^* . Let \mathbb{U} be the subset of $\mathbb{P}(\vec{P}^*, {}^{<\omega}\omega)$ consisting of conditions f_X such that for all $s \in X$ on level n , $f_X(s) \in \mathbb{U}_n$.

Proposition 6.9. *\mathbb{U} is dense in $\mathbb{P}(\vec{P}^*, {}^{<\omega}\omega)$.*

Next, we will prove the analogue of Proposition 2.5.

Lemma 6.10. *Every maximal antichain of $\mathbb{P}(\vec{P}, {}^{<\omega}\omega)$ from M remains maximal in $\mathbb{P}(\vec{P}^*, {}^{<\omega}\omega)$.*

Proof. Fix a maximal antichain $\mathcal{A} \in M$ of $\mathbb{P}(\vec{P}, {}^{<\omega}\omega)$. By Proposition 6.9, it suffices to show that every condition $f_X \in \mathbb{U}$ is compatible with some element of \mathcal{A} . So fix $f_X \in \mathbb{U}$. For a node $s \in X$ on level n of ${}^{<\omega}\omega$, let

$$f_X(s) = (\mathcal{T}_{j_0,s}^0 \wedge S_{0,s}, \dots, \dot{T}_{j_{n-1},s}^{n-1} \wedge \dot{S}_{n-1,s}).$$

Let $\langle j_0,s, \dots, j_{n-1},s \rangle = \vec{j}_s$, and note that we can have $\vec{j}_s = \vec{j}_{s'}$ for $s' \neq s$. Fix a condition $g_Y \in G$ forcing that $f_X \in \mathbb{U}$. Repeatedly using the construction in the proof of Lemma 6.7 on terminal nodes, we can find a determined condition $\bar{g}_Y \leq g_Y$ with $\bar{g}_Y(\vec{j}_s) = (\bar{q}_{\vec{j}_s}, \bar{F}_{\vec{j}_s})$ such that for every node $s \in X$, some $\tau_s : n \rightarrow {}^{<\omega}2$ lies on $(\bar{q}_{\vec{j}_s}, \bar{F}_{\vec{j}_s})$ with

$$\bar{q}_{\vec{j}_s}(0)_{\tau_s(0)} \leq S_{0,s}, \text{ and for all } 1 \leq i < n, (\bar{q}_{\vec{j}_s} \upharpoonright i) \upharpoonright (\tau_s \upharpoonright i) \Vdash \bar{q}_{\vec{j}_s}(i)_{\tau_s(i)} \leq \dot{S}_{i,s}.$$

The construction gives τ_s satisfying that if s' extends s , then $\tau_{s'}$ extends τ_s .

Let \bar{f}_X be a condition in $\mathbb{P}(\bar{P}, <^\omega \omega)$ such that for every node $s \in X$, we have $\bar{f}_X(s) = \bar{q}_{\bar{j}_s} \mid \tau_s$. Since \mathcal{A} is maximal in $\mathbb{P}(\bar{P}, <^\omega \omega)$, there is $a_Z \in \mathcal{A}$ compatible with \bar{f}_X . So let $r_W \leq a_Z, \bar{f}_X$.

We can carry out the construction in the proof of Proposition 4.8 to obtain a condition $h_Y \leq \bar{g}_Y$ with $h_Y(\bar{j}_s) = (q_{\bar{j}_s}, F_{\bar{j}_s})$ such that for every $s \in W$, some σ lies on $(q_{\bar{j}_s}, F_{\bar{j}_s})$ having $q_{\bar{j}_s} \mid \sigma \leq r_W(s)$. The condition h_Y forces that f_X and a_Z are compatible. By density, some such h_Y must be in G . \square

We now summarize our results in the following lemma, which will serve as the analogue of Proposition 2.4.

Lemma 6.11. *Suppose M is a suitable model and $\bar{P} = \langle \mathbb{P}_n \mid n < \omega \rangle \in M$ is an ω -iteration of perfect posets. If $G \subseteq \mathbb{Q}(\bar{P}, <^\omega \omega)$ is M -generic and $\bar{P}^* = \langle \mathbb{P}_n^* \mid n < \omega \rangle$ is constructed in $M[G]$ as above, then*

- (1) \bar{P}^* is an ω -iteration of perfect posets,
- (2) $\mathbb{Q}_0 \subseteq \mathbb{Q}_0^*$ and for all $n < \omega$, $\mathbb{1}_{\mathbb{P}_n^*}$ forces that $\dot{\mathbb{Q}}_n$ is a perfect poset and $\dot{\mathbb{Q}}_n^*$ extends it,
- (3) $\mathbb{P}_n \subseteq \mathbb{P}_n^*$,
- (4) every maximal antichain of \mathbb{P}_n from M remains maximal in \mathbb{P}_n^* ,
- (5) every maximal antichain of $\mathbb{P}(\bar{P}, <^\omega \omega)$ from M remains maximal in $\mathbb{P}(\bar{P}^*, <^\omega \omega)$.

Now we will define an ω -iteration $\bar{P}^J = \langle \mathbb{P}_n^J \mid n < \omega \rangle$ of perfect posets such that the tree iteration $\mathbb{P}(\bar{P}^J, <^\omega \omega)$ is going to have the property that in a forcing extension $L[G]$ of L by $\mathbb{P}(\bar{P}^J, <^\omega \omega)$, the only L -generic filters for the iteration \mathbb{P}_n^J are the restrictions of G to a level n node. This will be the generalization of Jensen's uniqueness of generic filters property to tree iterations.

Let $\langle S_\alpha \mid \alpha < \omega_1 \rangle$ be a canonically defined \diamond -sequence. We will construct \bar{P}^J in ω_1 -many steps using \diamond to seal maximal antichains along the way. Let \bar{P}_0 be the ω -iteration of perfect posets where $\mathbb{Q}_0 = \mathbb{P}_{\min}$ and each $\dot{\mathbb{Q}}_n = \check{\mathbb{P}}_{\min}$. Suppose the ω -iteration $\bar{P}_\alpha = \langle \mathbb{P}_n^\alpha \mid n < \omega \rangle$ of perfect poset has been defined. We let $\bar{P}_\alpha = \bar{P}_{\alpha+1}$, unless the following happens. Suppose S_α codes a well-founded binary relation $E \subseteq \alpha \times \alpha$ such that the collapse of E is a suitable model M_α with $\bar{P}_\alpha \in M_\alpha$ and $\alpha = \omega_1^{M_\alpha}$. In this case, we take the L -least M_α -generic filter $G \subseteq \mathbb{Q}(\bar{P}_\alpha, <^\omega \omega)$ and let $\bar{P}_{\alpha+1} = \bar{P}_\alpha^*$ as constructed in $M_\alpha[G]$. At limit stages λ , to obtain the ω -iteration \bar{P}_λ , we let \mathbb{Q}_0^λ be the union of the \mathbb{Q}_0^ξ for $\xi < \lambda$, and given that we have defined \mathbb{P}_n^λ , we let $\dot{\mathbb{Q}}_n^\lambda$ be a \mathbb{P}_n^λ -name for the poset that is the union of the $\dot{\mathbb{Q}}_n^\xi$ for $\xi < \lambda$. In order for this limit definition to make sense, we need to verify that each $\dot{\mathbb{Q}}_n^\xi$ is a \mathbb{P}_n^λ -name for a perfect poset. So let's argue that this is indeed the case.

Clearly \mathbb{P}_1^λ makes sense, $\mathbb{P}_1^\xi \subseteq \mathbb{P}_1^\lambda$ for every $\xi < \lambda$, and every maximal antichain of \mathbb{P}_1^ξ from M_ξ remains maximal in \mathbb{P}_1^λ . So we can assume inductively that we have defined \mathbb{P}_n^λ so that $\mathbb{P}_n^\xi \subseteq \mathbb{P}_n^\lambda$ for every $\xi < \lambda$, and every maximal antichain of \mathbb{P}_n^ξ in M_ξ remains maximal in \mathbb{P}_n^λ . Let $H^* \subseteq \mathbb{P}_n^\lambda$ be V -generic. By our assumption, the filter H^* restricts to an M_ξ -generic filter H for \mathbb{P}_n^ξ . The model $M_\xi[H]$ satisfies that $(\dot{\mathbb{Q}}_n^\xi)_H$ is a perfect poset and so this must be the case in $V[H^*]$ as well since this is absolute. So we can extend the definition to \mathbb{P}_{n+1}^λ . It is easy to see that $\mathbb{P}_{n+1}^\xi \subseteq \mathbb{P}_{n+1}^\lambda$. It remains to argue that every maximal antichain $\mathcal{A} \in M_\xi$ of \mathbb{P}_{n+1}^ξ remains maximal in \mathbb{P}_{n+1}^λ . Fix $p \in \mathbb{P}_{n+1}^\lambda$. We can assume inductively that we have

already showed for every $\mu < \nu < \lambda$, that every maximal antichain of \mathbb{P}_{n+1}^μ in M_μ remains maximal in \mathbb{P}_{n+1}^ν . Then, by definition of \mathbb{P}_{n+1}^λ , there is some $\xi < \eta < \lambda$ such that $p(0) \in \mathbb{Q}_0^\eta$, and $p \upharpoonright i \Vdash p(i) \in \dot{\mathbb{Q}}_i^\eta$ for $1 \leq i < n$. Since \mathcal{A} is maximal in \mathbb{P}_{n+1}^η , it follows that there is some $q \in \mathcal{A}$ that is compatible with $p \in \mathbb{P}_{n+1}^\eta \subseteq \mathbb{P}_{n+1}^\lambda$. In particular, we have just shown the following.

Lemma 6.12. *If λ is a limit ordinal and $\xi < \lambda$, then $\mathbb{P}_n^\xi \subseteq \mathbb{P}_n^\lambda$ and every maximal antichain of \mathbb{P}_n^ξ from M_ξ remains maximal in \mathbb{P}_n^λ .*

Lemma 6.13. *If $\alpha < \beta < \omega_1$, then every maximal antichain of $\mathbb{P}(\vec{P}_\alpha, <^\omega \omega)$ from M_α remains maximal in $\mathbb{P}(\vec{P}_\beta, <^\omega \omega)$.*

The proof uses Lemma 6.8 for successor stages and follows easily for limit stages.

For $n < \omega$, let $\mathbb{P}_n^J = \mathbb{P}_n^{\omega_1}$ be constructed as all the limit stages and let $\vec{P}^J = \langle \mathbb{P}_n^J \mid n < \omega \rangle$ be the corresponding ω -iteration.

It will be useful for future arguments to assume that conditions in \mathbb{P}_n^J are coded by reals.

Theorem 6.14. *The poset $\mathbb{P}(\vec{P}^J, <^\omega \omega)$ has the ccc.*

Proof. Fix a maximal antichain \mathcal{A} of $\mathbb{P}(\vec{P}^J, <^\omega \omega)$. Choose some transitive $M \prec L_{\omega_2}$ of size ω_1 with $\mathcal{A} \in M$. We can decompose M as the union of an elementary chain of countable substructures

$$X_0 \prec X_1 \prec \cdots \prec X_\alpha \prec \cdots \prec M$$

with $\mathcal{A} \in X_0$. By the properties of \diamond , there is some α such that $\alpha = \omega_1 \cap X_\alpha$, $\mathbb{P}_n^\alpha = \mathbb{P}_n^J \cap X_\alpha$ for all $n < \omega$, and S_α codes X_α . Let M_α be the transitive collapse of X_α . Then \mathbb{P}_n^α is the image of \mathbb{P}_n^J under the collapse and α is the image of ω_1 . Let $\bar{\mathcal{A}} = \mathcal{A} \cap X_\alpha$ be the image of \mathcal{A} . So at stage α in the construction of \vec{P}^J , we chose a forcing extension $M_\alpha[G]$ of M_α by $\mathbb{Q}(\vec{P}_\alpha, <^\omega \omega)$ and let $\mathbb{P}_n^{\alpha+1} = (\mathbb{P}_n^\alpha)^*$ be constructed in $M_\alpha[G]$. By Lemma 6.13, $\bar{\mathcal{A}}$ remains maximal in all further $\mathbb{P}(\vec{P}_\beta, <^\omega \omega)$ for $\beta > \alpha$, and hence $\bar{\mathcal{A}}$ remains maximal in $\mathbb{P}(\vec{P}^J, <^\omega \omega)$. It follows that $\bar{\mathcal{A}} = \mathcal{A}$ is countable. \square

7. THE GENERALIZED KANOVEI-LYUBETSKY THEOREM

Now that we have constructed the ω -iteration $\vec{P}^J = \langle \mathbb{P}_n^J \mid n < \omega \rangle$ and the corresponding tree iteration $\mathbb{P}(\vec{P}^J, <^\omega \omega)$ of Jensen's forcing, we would like to generalize the Kanovei-Lyubetsky argument of Section 3 to prove the analogue of the “uniqueness of generics” property for tree iterations. Namely, we will show that in a forcing extension $L[G]$ by $\mathbb{P}(\vec{P}^J, <^\omega \omega)$ the only L -generic filters for \mathbb{P}_n^J are the restrictions of G to a level n node.

Suppose that $\vec{P} = \langle \mathbb{P}_n \mid n < \omega \rangle$ is an ω -iteration of perfect posets and G is a generic filter for $\mathbb{P}(\vec{P}, <^\omega \omega)$. Given a node s on level n of $<^\omega \omega$, let x_s be the n -length sequence of generic reals added by G on node s and let \dot{x}_s be the canonical name for x_s .

For the next lemma, suppose that $\vec{P} = \langle \mathbb{P}_n \mid n < \omega \rangle$ is an ω -iteration of perfect posets that is an element of a suitable model M . We should think of \vec{P} as one of the ω -iterations \vec{P}_α arising at stage α in the construction of the ω -iteration \vec{P}^J and we should think of M as the model M_α from that stage.

Lemma 7.1. *In M , suppose that \dot{r} is a $\mathbb{P}(\vec{P}, <^\omega\omega)$ -name for an n -length sequence of reals such that for all nodes s on level n of $<^\omega\omega$,*

$$\mathbb{1} \Vdash_{\mathbb{P}(\vec{P}, <^\omega\omega)} \dot{r} \neq \dot{x}_s.$$

Then in a forcing extension $M[G]$ by $\mathbb{Q}(\vec{P}, <^\omega\omega)$, for every node s on level n of $<^\omega\omega$, the set of conditions forcing the statement

$$\Phi(s) := \text{“If } \dot{r} \text{ is (ground model) generic for } \mathbb{P}_n^*, \text{ then } \langle \mathcal{T}_{s(0)}^0, \mathcal{T}_{s(1)}^1, \dots, \mathcal{T}_{s(n-1)}^{n-1} \rangle \text{ is not in the filter determined by } \dot{r} \text{.”}$$

is dense in $\mathbb{P}(\vec{P}^, <^\omega\omega)$.*

Proof. Fix a condition $f_X \in \mathbb{P}(\vec{P}^*, <^\omega\omega)$ and a node d on level n of $<^\omega\omega$. We need to find a condition $f'_X \leq f_X$ such that $f'_X \Vdash \Phi(d)$. Since \mathbb{U} is dense in $\mathbb{P}(\vec{P}^*, <^\omega\omega)$, we can assume without loss that $f_X \in \mathbb{U}$. For a node s on level m of X , let

$$f_X(s) = (\mathcal{T}_{j_{0,s}}^0 \wedge S_{0,s}, \dots, \mathcal{T}_{j_{m-1,s}}^{m-1} \wedge \dot{S}_{m-1,s}).$$

Let $\langle j_{0,s}, \dots, j_{m-1,s} \rangle = \vec{j}_s$, and note that we can have $\vec{j}_s = \vec{j}_{s'}$. By strengthening if necessary, we can assume that there is a node $s \in X$ with $\vec{j}_s = d$. Repeatedly using the construction in the proof of Lemma 6.7 on terminal nodes, we can find a determined condition $g_Y \in G$ with $g_Y(\vec{j}_s) = (q_{\vec{j}_s}^{\vec{j}_s}, F_{\vec{j}_s}^{\vec{j}_s})$ such that for every node $s \in X$, some $\tau_s : m \rightarrow <^\omega 2$ lies on $(q_{\vec{j}_s}^{\vec{j}_s}, F_{\vec{j}_s}^{\vec{j}_s})$ with

$$q_{\vec{j}_s}^{\vec{j}_s}(0)_{\tau_s(0)} \leq S_{0,s}, \text{ and for all } 1 \leq i < m, (q_{\vec{j}_s}^{\vec{j}_s} \upharpoonright i) \mid (\tau_s \upharpoonright i) \Vdash q_{\vec{j}_s}^{\vec{j}_s}(i)_{\tau_s(i)} \leq \dot{S}_{i,s}.$$

The construction gives τ_s satisfying that if s' extends s , then $\tau_{s'}$ extends τ_s .

Now we are going to construct a condition $a_A^g \in \mathbb{P}(\vec{P}, <^\omega\omega)$ with $a_A^g(s) = q_s$ associated to g_Y , satisfying the following properties:

- (1) $X \subseteq A$ and for every node $s \in X$, $q_s \leq q_{\vec{j}_s}^{\vec{j}_s} \mid \tau_s$.
There is $X \subseteq \bar{A} \subseteq A$ such that:
- (2) For every σ which lies on (q_d, F_d) , there is a node $s_\sigma \in \bar{A}$ such that $q_{s_\sigma} \leq q_d \mid \sigma$.
- (3) For every node s on level n of \bar{A} , a_A^g forces over $\mathbb{P}(\vec{P}, <^\omega\omega)$ the statement:
“There is $i < n - 1$ such that $\dot{r} \upharpoonright i$ is (ground model) generic for \mathbb{P}_i and $\dot{r}(i) \notin [(a_A^g(s)(i))_{\dot{r} \upharpoonright i}]$.”

Let $a_{\bar{A}}$ be a condition in $\mathbb{P}(\vec{P}, <^\omega\omega)$ with $X \subseteq \bar{A}$ be such that $a_{\bar{A}}(s) = q_{\vec{j}_s}^{\vec{j}_s} \mid \tau_s$ for every node $s \in X$, and for every σ which lies on (q_d, F_d) , there is a node $s_\sigma \in \bar{A}$ such that $a_{\bar{A}}(s_\sigma) = q_d \mid \sigma$. Fix a node s on level n of \bar{A} and consider a forcing extension $M[H]$ by $\mathbb{P}(\vec{P}, <^\omega\omega)$. By assumption, we have $r = \dot{r}_H \neq x_s$. So there is $i < n - 1$ such that $r \upharpoonright i = x_s \upharpoonright i$ and $r(i) \neq x_s(i)$. So we can strengthen $a_{\bar{A}}$ to a condition $a_{A'}$ forcing that there is $i < n - 1$ such that $\dot{r} \upharpoonright i$ is M -generic and $\dot{r}(i) \notin [(a_{A'}(s)(i))_{\dot{r} \upharpoonright i}]$. By repeating this for all the finitely many nodes on level n , we obtain the required condition a_A^g .

Next, we can carry out the construction in the proof of Proposition 4.8 to obtain a condition $\bar{g}_Y \leq g_Y$ with $\bar{g}_Y(\vec{j}_s) = (\bar{q}_{\vec{j}_s}^{\vec{j}_s}, \bar{F}_{\vec{j}_s}^{\vec{j}_s})$ such that for every $s \in X$, some σ lies on $(\bar{q}_{\vec{j}_s}^{\vec{j}_s}, \bar{F}_{\vec{j}_s}^{\vec{j}_s})$ having $\bar{q}_{\vec{j}_s}^{\vec{j}_s} \mid \sigma \leq a_A^g(s)$. We can also ensure that for every τ which lies on (\bar{q}_d, \bar{F}_d) , there is some σ which lies on (q_d, F_d) such that τ extends σ and $\bar{q}_d \mid \tau \leq a_A^g(s_\sigma)$. By density, some such \bar{g}_Y must be in G .

By construction, \bar{g}_Y forces that f_X is compatible with a_A^g . So let $\bar{f}_X \leq f_X, a_A^g$. We will be done if we can show that a_A^g forces the statement $\Phi(d)$ over $\mathbb{P}(\vec{P}^*, <^\omega\omega)$.

Suppose $H^* \subseteq \mathbb{P}(\vec{P}^*, <^\omega\omega)$ is a V -generic filter containing a_A^g ($V = M[G]$). Now let's suppose towards a contradiction that $p = \langle \mathcal{T}_{d(0)}^0, \dot{\mathcal{T}}_{d(1)}^1, \dots, \dot{\mathcal{T}}_{d(n-1)}^{n-1} \rangle$ is in the filter determined by r . Thus, there is some σ which lies on p such for all $i < n$, $r(i)$ is a branch through $(p \upharpoonright \sigma(i))_{r \upharpoonright i}$. By construction, we have that $p \upharpoonright \sigma \leq a_A^g(s)$ for some s on level n of $<^\omega\omega$. Let H be the restriction of H^* to an M -generic filter for $\mathbb{P}(\vec{P}, <^\omega\omega)$, and note that $a_A^g \in H$. Thus, there is some $i < n - 1$ such that $r(i)$ is not a branch through $(a_A^g(s)(i))_{r \upharpoonright i}$, which is the desired contradiction. \square

Theorem 7.2. *Suppose $H \subseteq \mathbb{P}(\vec{P}^J, <^\omega\omega)$ is L -generic. If an n -length sequence of reals $r \in L[H]$ is L -generic for \mathbb{P}_n , then $r = x_s$ for some node s on level n of $<^\omega\omega$.*

Proof. Let's suppose that r is not one of the x_s for s on level n of $<^\omega\omega$. Let \dot{r} be a nice $\mathbb{P}(\vec{P}^J, <^\omega\omega)$ -name for r such that for all nodes s on level n of $<^\omega\omega$, $\mathbb{1} \Vdash_{\mathbb{P}(\vec{P}^J, <^\omega\omega)} \dot{r} \neq \dot{x}_s$.

Choose some transitive $N \prec L_{\omega_2}$ of size ω_1 with $\dot{r} \in N$. We can decompose N as the union of a continuous elementary chain of countable substructures

$$X_0 \prec X_1 \prec \dots \prec X_\alpha \prec \dots \prec N$$

with $\dot{r} \in X_0$. By the properties of \diamond , there is some α such that $\alpha = \omega_1 \cap X_\alpha$, $\mathbb{P}_n^\alpha = \mathbb{P}_n^J \cap X_\alpha$ for all $n < \omega$, and S_α codes X_α . Let M_α be the transitive collapse of X_α . Then, for every $n < \omega$, \mathbb{P}_n^α is the image of \mathbb{P}_n^J under the collapse, and α is the image of ω_1 . The name \dot{r} is fixed by the collapse by our assumption that we can always code conditions in $\mathbb{P}(\vec{P}^J, <^\omega\omega)$ by subsets of ω and because all antichains of $\mathbb{P}(\vec{P}^J, <^\omega\omega)$ are countable. So at stage α in the construction of \vec{P}^J , we chose a forcing extension $M_\alpha[G]$ of M_α by $\mathbb{Q}(\vec{P}^\alpha, <^\omega\omega)$ and let $\vec{P}_{\alpha+1} = \vec{P}_\alpha^*$ be constructed in $M_\alpha[G]$.

By elementarity, M_α satisfies that $\mathbb{1}_{\mathbb{P}(\vec{P}_\alpha, <^\omega\omega)} \Vdash \dot{r} \neq \dot{x}_s$ for all s on level n of $<^\omega\omega$. Thus, by Lemma 7.1, for every s on level n of $<^\omega\omega$, $\mathbb{P}(\vec{P}_{\alpha+1}, <^\omega\omega)$ has a maximal antichain \mathcal{A}_s consisting of conditions f_X forcing the statement:

$\Phi(s) :=$ "If \dot{r} is (ground model) generic for \mathbb{P}_n^* , then $\langle \mathcal{T}_{s(0)}^0, \dot{\mathcal{T}}_{s(1)}^1, \dots, \dot{\mathcal{T}}_{s(n-1)}^{n-1} \rangle$ is not in the filter determined by \dot{r} ."

It follows, by an argument analogous to the proof of Proposition 2.7, that every \mathcal{A}_s remains maximal in $\mathbb{P}(\vec{P}^J, <^\omega\omega)$. So let's argue that if $f_X \in \mathcal{A}_s$, then f_X forces $\Phi(s)$ over $\mathbb{P}(\vec{P}^J, <^\omega\omega)$.

Let $\bar{H}^* \subseteq \mathbb{P}(\vec{P}^J, <^\omega\omega)$ be an L -generic filter containing f_X and let \bar{H} be the restriction of \bar{H}^* to $\mathbb{P}(\vec{P}_{\alpha+1}, <^\omega\omega)$. Let $r = \dot{r}_{\bar{H}^*}$. Since $f_X \in \bar{H}$, it follows that $M_\alpha[G_\alpha][\bar{H}]$ satisfies that if r is $M_\alpha[G_\alpha]$ -generic, then $\langle \mathcal{T}_{s(0)}^0, \dot{\mathcal{T}}_{s(1)}^1, \dots, \dot{\mathcal{T}}_{s(n-1)}^{n-1} \rangle$ is not in the filter determined by r . If r is in fact fully L -generic, then the statement will continue to hold in $L[\bar{H}^*]$.

Since H must meet every \mathcal{A}_s , it holds in $L[H]$ that if r is L -generic for \mathbb{P}_n^J , then it does not meet the maximal antichain consisting of conditions $\langle \mathcal{T}_{s(0)}^0, \dot{\mathcal{T}}_{s(1)}^1, \dots, \dot{\mathcal{T}}_{s(n-1)}^{n-1} \rangle$ for nodes s on level n of $<^\omega\omega$ (from level α of the construction), and so r in fact cannot be L -generic. \square

8. ITERATING \vec{P}^J ALONG THE TREE ${}^{<\omega}\omega_1$

We will now argue that the tree iteration $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$, where we iterate along the uncountable tree ${}^{<\omega}\omega_1$, shares all the key properties of the tree iteration $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega)$, namely it has the ccc and Theorem 7.2, concerning the uniqueness of generic filters for \mathbb{P}_n^J , continues to hold.

Theorem 8.1. *The poset $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$ has the ccc.*

Proof. Let's suppose to the contrary that there is an uncountable antichain \mathcal{A} in $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$. By a Δ -system argument, there must be some finite subtree $X \subseteq {}^{<\omega}\omega_1$ and an uncountable subset $\mathcal{A}' \subseteq \mathcal{A}$ such that for any f_Y and g_Z in \mathcal{A}' , $Y \cap Z = X$. Given $f_Y \in \mathcal{A}'$, let f_X be the restriction of f_Y to X . Clearly the collection of all such f_X is an uncountable antichain in $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$. But then since X is finite, there must be a corresponding uncountable antichain in $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega)$, which is impossible by Theorem 6.14. \square

Theorem 8.2. *Suppose $H \subseteq \mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$ is L -generic. If an n -length sequence of reals $r \in L[H]$ is L -generic for \mathbb{P}_n^J , then $r = x_s$ for some node s on level n of ${}^{<\omega}\omega_1$.*

Proof. Suppose that r is an n -length sequence of reals in $L[H]$ that is L -generic for \mathbb{P}_n^J . Let \dot{r} be a nice $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$ -name for r . Since $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$ has the ccc by Theorem 8.2, it follows that conditions in the name \dot{r} use only countably many nodes of ${}^{<\omega}\omega_1$. Thus, \dot{r} is a $\mathbb{P}(\vec{P}, \mathcal{T})$ -name, where \mathcal{T} is a countable subtree of ${}^{<\omega}\omega_1$. We may assume that \mathcal{T} is isomorphic to ${}^{<\omega}\omega$, and so $\mathbb{P}(\vec{P}, \mathcal{T})$ is isomorphic to $\mathbb{P}(\vec{P}, {}^{<\omega}\omega)$. Let \bar{H} be the restriction of H to $\mathbb{P}(\vec{P}, \mathcal{T})$, which is L -generic for it. Thus, $r \in L[\bar{H}]$, from which it follows, by Theorem 7.2, that $r = x_s$ for some $s \in \mathcal{T}$. \square

9. SYMMETRIC MODELS OF $ZF + AC_\omega + \neg DC$

We will construct a symmetric submodel of a forcing extension $L[G]$ by $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$ in which $ZF + AC_\omega$ holds, but DC fails. The reals of this model will be a model of second-order arithmetic in which $Z_2 + \Sigma_\infty^1$ -AC holds, but Π_2^1 -DC fails. Let's start with a brief discussion of the method of constructing symmetric submodels of a forcing extension, which goes all the way back to Cohen's pioneering work on forcing.

Suppose that \mathbb{P} is a forcing notion. Recall that if π is an automorphism of \mathbb{P} , then we can apply π recursively to conditions in a \mathbb{P} -name σ to obtain the \mathbb{P} -name $\pi(\sigma)$. It is not difficult to see by induction on complexity of formulas, that for every formula φ and condition $p \in \mathbb{P}$, $p \Vdash \varphi(\sigma)$ if and only if $\pi(p) \Vdash \varphi(\pi(\sigma))$. Fix some group \mathcal{G} of automorphisms of \mathbb{P} . Recall that a filter \mathcal{F} on subgroups of a group \mathcal{G} is *normal* if whenever $g \in \mathcal{G}$ and $\mathcal{K} \in \mathcal{F}$, then $g\mathcal{K}g^{-1} \in \mathcal{F}$. Let's fix a normal filter \mathcal{F} on the subgroups of \mathcal{G} . The subgroup of \mathcal{G} fixing a particular \mathbb{P} -name σ , consisting of automorphisms π such that $\pi(\sigma) = \sigma$, is called $\text{sym}(\sigma)$. If $\text{sym}(\sigma)$ is in \mathcal{F} , then we say that σ is a *symmetric* \mathbb{P} -name. We recursively define that a \mathbb{P} -name is *hereditarily symmetric* when it is symmetric and all names inside it are hereditarily symmetric. Let HS be the collection of all hereditarily symmetric \mathbb{P} -names. Let $G \subseteq \mathbb{P}$ be V -generic. The *symmetric model*

$$N = \{\sigma_G \mid \sigma \in \text{HS}\}$$

associated to the group of automorphisms \mathcal{G} and the normal filter \mathcal{F} consists of the interpretations of all hereditarily symmetric \mathbb{P} -names. It is a standard result that $N \models \text{ZF}$.

We would like to review now a classical construction of a symmetric model in which countable choice holds but dependent choice fails, which is due to Jensen [Jen67]. Let \mathbb{P} be the forcing adding a set of Cohen subsets of ω_1 indexed by the tree ${}^{<\omega}\omega_1$ with countable conditions. We will call $\text{Add}(\omega_1, 1)$ the poset adding a Cohen subset to ω_1 with countable conditions. So, more precisely, conditions in \mathbb{P} are functions $f_X : X \rightarrow \text{Add}(\omega_1, 1)$, where X is some countable subtree of ${}^{<\omega}\omega_1$, ordered so that $g_Y \leq f_X$ whenever $X \subseteq Y$ and for all $t \in X$, $g_Y(t) \leq f_X(t)$ in $\text{Add}(\omega_1, 1)$. Note that \mathbb{P} is countably closed. Next, we need to select an appropriate group of automorphisms of \mathbb{P} .

Let $\text{Aut}({}^{<\omega}\omega_1)$ be the automorphism group of the tree ${}^{<\omega}\omega_1$. Every automorphism $\pi \in \text{Aut}({}^{<\omega}\omega_1)$ induces an automorphism π^* of \mathbb{P} , so that $\pi^*(f_X) = f_{\pi \smallfrown X}$ with $f_{\pi \smallfrown X}(t) = f_X(\pi^{-1}(t))$. Let

$$\mathcal{G} = \{\pi^* \mid \pi \in \text{Aut}({}^{<\omega}\omega_1)\}.$$

Next, we need to select an appropriate filter on the subgroups of \mathcal{G} . Let's call a subtree T of the tree ${}^{<\omega}\omega_1$ *useful* if it is countable and doesn't have an infinite branch. Given a useful tree T , let H_T be the subgroup of \mathcal{G} consisting of all automorphisms π^* such that π point-wise fixes T . Let \mathcal{F} be the filter on the subgroups of \mathcal{G} generated by all such subgroups H_T . To see that \mathcal{F} is normal, observe that if T is a useful tree and $H_T \subseteq K \in \mathcal{F}$, then $\pi \smallfrown T$ is useful and $H_{\pi \smallfrown T} \subseteq \pi^* K \pi^{*-1}$.

Now, let $G \subseteq \mathbb{P}$ be V -generic and let $N = \{\sigma_G \mid \sigma \in \text{HS}\}$. In $V[G]$, let \mathcal{T} be the tree isomorphic to ${}^{<\omega}\omega_1$ whose nodes are the Cohen subsets of ω_1 added by G . Given a node $t \in {}^{<\omega}\omega_1$, let σ_t be the canonical name for the Cohen subset of ω_1 added on node t by G . Let $\dot{\mathcal{T}} = \{(\text{op}(\sigma_s, \sigma_t), \mathbb{1}_{\mathbb{P}}) \mid s \leq t \text{ in } {}^{<\omega}\omega_1\}$, where $\text{op}(\sigma_s, \sigma_t)$ is the canonical \mathbb{P} -name for the ordered pair of σ_s and σ_t . Clearly $\dot{\mathcal{T}}_G = \mathcal{T}$. Fix any $\pi^* \in \mathcal{G}$, and observe that $\pi^*(\dot{\mathcal{T}}) = \{(\text{op}(\sigma_{\pi(s)}, \sigma_{\pi(t)}), \mathbb{1}_{\mathbb{P}}) \mid s \leq t \text{ in } {}^{<\omega}\omega_1\} = \dot{\mathcal{T}}$. Also, any automorphism π^* with $\pi(s) = s$ fixes σ_s . This shows that $\dot{\mathcal{T}} \in \text{HS}$, and hence \mathcal{T} is in the symmetric model N . Using that a symmetric name must be fixed by a group of automorphisms point-wise fixing a tree without an infinite branch, we can show that no infinite branch through \mathcal{T} can have a symmetric name, and so DC fails in N . Using that \mathbb{P} is countably closed, we can show that AC_ω holds in N .

We are now ready to construct a symmetric model N of a forcing extension $L[G]$ by $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$ with the same properties. First, observe that every $\pi \in \text{Aut}({}^{<\omega}\omega_1)$ induces an automorphism π^* of $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$ so that $\pi^*(f_X) = f_{\pi \smallfrown X}$ with $f_{\pi \smallfrown X}(t) = f_X(\pi^{-1}(t))$. As above, let \mathcal{G} be the group of automorphisms π^* for $\pi \in \text{Aut}({}^{<\omega}\omega_1)$ and let \mathcal{F} be the normal filter generated by the subgroups H_T of \mathcal{G} , consisting of all automorphisms point-wise fixing some useful subtree T of $\omega_1^{<\omega}$. Note that in the present situation the domains of conditions f_X are finite subtrees X , while we use countable trees T to determine which names are symmetric. Let $N \subseteq L[G]$ be the symmetric model determined by the group of automorphisms \mathcal{G} and the filter \mathcal{F} . In $L[G]$, consider the tree \mathcal{T} , isomorphic to ${}^{<\omega}\omega_1$ whose nodes are the generic sequences of reals added by G . The argument we gave above generalizes

in a straightforward way to show that $\dot{\mathcal{T}}$, the canonical name for \mathcal{T} , is hereditarily symmetric, and hence $\mathcal{T} \in N$.

Suppose T is any subtree of $\omega_1^{<\omega}$. If f_X is a condition in $\mathbb{P}(\vec{P}^J, <^\omega\omega_1)$, we will denote by $f_{X \cap T}$, the restriction of f_X to nodes in T , i.e. $f_{X \cap T}$ has domain $X \cap T$ and $f_{X \cap T}(t) = f_X(t)$. We let

$$G_T = \{f_X \in G \mid X \subseteq T\}.$$

Note that if T is useful, then the canonical name for G_T is fixed by all elements of the subgroup H_T , and therefore $G_T \in N$. Thus, each $L[G_T] \subseteq N$.

If σ is a $\mathbb{P}(\vec{P}^J, <^\omega\omega_1)$ -name and $H_T \subseteq \text{sym}(\sigma)$ for a useful tree T , we will say that T *witnesses* that σ is symmetric.

Proposition 9.1. *Suppose that σ is a symmetric $\mathbb{P}(\vec{P}^J, <^\omega\omega_1)$ -name, as witnessed by a useful tree T , and $f_X \Vdash \varphi(\sigma, \check{a})$. Then $f_{X \cap T} \Vdash \varphi(\sigma, \check{a})$.*

Proof. Suppose it is not the case that $f_{X \cap T} \Vdash \varphi(\sigma, \check{a})$. Then there is a condition $g_Y \leq f_{X \cap T}$ which forces $\neg\varphi(\sigma, \check{a})$. So g_Y is incompatible with f_X and the incompatibility must occur on nodes outside of T . Let $\pi \in \text{Aut}(<^\omega\omega_1)$ be an automorphism which switches these nodes in $Y \setminus T$ with some nodes outside both T and X . Clearly $\pi^* \in H_T$, and so $\pi^*(\sigma) = \sigma$. It follows that $\pi^*(g_Y) \Vdash \neg\varphi(\sigma, \check{a})$. But, by construction, $\pi^*(g_Y)$ is compatible with f_X , which is the desired contradiction. Thus, we have shown that $f_{X \cap T} \Vdash \varphi(\sigma, \check{a})$. \square

Proposition 9.2. *Suppose that σ is a symmetric $\mathbb{P}(\vec{P}^J, <^\omega\omega_1)$ -name, as witnessed by a useful tree T , and $A = \sigma_G$ is a set of ordinals. Then $A \in L[G_T]$.*

Proof. Let f_X be some condition in G forcing that σ is a set of ordinals. Define a name

$$\sigma^* = \{(\xi, g_{Y \cap T}) \mid g_Y \leq f_X, g_Y \Vdash \xi \in \sigma\}.$$

We will argue that $f_X \Vdash \sigma = \sigma^*$. Let $H \subseteq \mathbb{P}(\vec{P}^J, <^\omega\omega_1)$ be some L -generic filter containing f_X . Suppose $\xi \in \sigma_H$. Then there is $g_Y \in H$ such that $g_Y \leq f_X$ and $g_Y \Vdash \xi \in \sigma$, from which it follows that $(\xi, g_{Y \cap T}) \in \sigma^*$ and $g_{Y \cap T} \in H$. So we have $\xi \in \sigma_H^*$. Next, suppose that $\xi \in \sigma_H^*$. Then there is a condition $g_Y \Vdash \xi \in \sigma$ such that $(\xi, g_{Y \cap T}) \in \sigma^*$ and $g_{Y \cap T} \in H$. But by Proposition 9.1, it follows that $g_{Y \cap T} \Vdash \xi \in \sigma$, and so $\xi \in \sigma_H$. \square

Lemma 9.3. *DC fails in N .*

Proof. We will argue that \mathcal{T} does not have an infinite branch in N , and hence DC fails. Suppose to the contrary that N has an infinite branch b through \mathcal{T} , which we can view via coding as a subset of natural numbers (since it is an ω -length sequence of reals). Fix a symmetric name \dot{b} for b , as witnessed by a useful tree T . By Proposition 9.2, we can assume that the name \dot{b} mentions only conditions with domains contained in T . Recall that for a node $s \in <^\omega\omega_1$, x_s is the L -generic sequence of reals for $\mathbb{P}_{\text{len}(s)}^J$ added by G on node s and \dot{x}_s is the canonical $\mathbb{P}(\vec{P}^J, <^\omega\omega_1)$ -name for x_s . Let \bar{b} be the branch through the tree $<^\omega\omega_1$ which corresponds to b via the obvious isomorphism. Since T doesn't have infinite branches by assumption, \bar{b} cannot be a branch through T . Thus, there is some natural number n such that $\bar{b} \upharpoonright n \subseteq T$ and $\bar{b}(n) = s$ is outside T . Fix a condition

$$f_X \Vdash \dot{b}(n) = \dot{x}_s.$$

Let π be any automorphism of the tree ${}^{<\omega}\omega_1$ fixing $T \cup X$ which swaps s with some node t . By genericity, $x_s \neq x_t$. Since π fixes $T \cup X$, it follows that $\pi^*(\dot{b}) = \dot{b}$ and $\pi^*(f_X) = f_X$. Let $H = \pi \restriction G$, which is also L -generic for $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$. Observe that $\dot{b}_H = \dot{b}$ since the name \dot{b} only mentioned conditions with domain in T and T is fixed by π . Since $f_X \in H$, it follows that $b(n) = (\dot{x}_s)_H$, but this is impossible since $(\dot{x}_s)_H = x_t$ and $(\dot{x}_s)_G = x_s$. \square

The proof of the next lemma relies mainly on the fact that $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$ has the ccc and uses very little else about what the conditions in $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$ look like. So to make the notation nicer, we will switch away from the previous convention and call conditions in $\mathbb{P}(\vec{P}^J, {}^{<\omega}\omega_1)$ standard names like p and q .

Lemma 9.4. *AC $_{\omega}$ holds in N .*

Proof. Suppose $F \in N$ is a countable family of non-empty sets. Let \dot{F} be a hereditarily symmetric name for F with the useful tree S witnessing that \dot{F} is symmetric, and let $q \in G$ force that \dot{F} is a countable family of non-empty sets. We would like to build a name $\dot{C} \in \text{HS}$ such that q forces that \dot{C} is a choice function for \dot{F} . We will adapt the following strategy. First, we will build a mixed name $\dot{C}_0 \in \text{HS}$ and a useful tree T_0 extending S , witnessing that \dot{C}_0 is symmetric, such that $q \Vdash \dot{C}_0 \in \dot{F}(0)$. Next, we will build a mixed name $\dot{C}_1 \in \text{HS}$ and a useful tree T_1 extending T_0 , witnessing that \dot{C}_1 is symmetric, such that $q \Vdash \dot{C}_1 \in \dot{F}(1)$. Proceeding in this fashion, we will build names $\dot{C}_n \in \text{HS}$ and an increasing sequence of useful trees T_n such that $q \Vdash \dot{C}_n \in \dot{F}(n)$. Provided we can ensure in the course of the construction that $T = \bigcup_{n \in \omega} T_n$ does not have an infinite branch, we will be able to build from the names \dot{C}_n a hereditarily symmetric name \dot{C} , witnessed by T to be symmetric, that is forced by q to be a choice function for \dot{F} .

Let D_0 be the dense set below q of conditions p such that for some name $\dot{c}_p \in \text{HS}$, $p \Vdash \dot{c}_p \in \dot{F}(0)$. We will thin out D_0 to a maximal antichain over which we can mix the names \dot{c}_p to get the desired name $\dot{C}_0 \in \text{HS}$. At the beginning, we choose some condition $p_0 \in D_0$ and a name $\dot{c}_{p_0} \in \text{HS}$, witnessed by a useful tree S'_0 to be symmetric, such that $p_0 \Vdash \dot{c}_{p_0} \in \dot{F}(0)$. By extending S'_0 , if necessary, we can assume that $\text{dom}(p_0) \subseteq S'_0$. Let S_0 be the union of S and S'_0 . Next, we choose some $p'_1 \in D_0$, incompatible to p_0 , and a name $\dot{c}_{p'_1} \in \text{HS}$, witnessed by a useful tree S'_1 with $\text{dom}(p'_1) \subseteq S'_1$ to be symmetric, such that $p'_1 \Vdash \dot{c}_{p'_1} \in \dot{F}(0)$. At this point, it is not a good idea to union up S_0 and S'_1 because if we keep doing this we might end up with an infinite branch in the final tree. So instead, we first consider those nodes in $S_0 \setminus S$ which have some new node of S'_1 sitting directly above them, meaning that we are extending an old branch that was already extended once from S_0 . Call these nodes in $S_0 \setminus S$ *bad* and the nodes of S'_1 sitting directly above them *bad successors*. Let π be an automorphism of the tree ${}^{<\omega}\omega_1$ which moves every bad node t (and its bad successors along with it) to some node outside S_0 . Since S_0 is countable and each node in ${}^{<\omega}\omega_1$ has ω_1 -many successors, there are many nodes to which the nodes t can move. Since π fixes S , we have $\pi^* \in H_S$. So we have $\pi^*(p'_1) \Vdash \pi^*(\dot{c}_{p'_1}) \in \dot{F}(0)$. We can safely union up S_0 and $\pi^*(S'_1)$ without growing any branches in S_0 that have extended beyond S . But the problem now is that $\pi^*(p'_1)$ may no longer be incompatible with p_0 , since the incompatibility could have come from some node t that got moved. This, however, can be easily remedied by adding back to $\pi^*(p'_1)$ one of the nodes from S_0 which caused the incompatibility. Let p_1 be the condition

$\pi^*(p'_1)$ together with possibly one extra node from $S'_1 \cap S_0$, let $\dot{c}_{p_1} = \pi^*(\dot{c}_{p'_1})$, and let S_1 be the union of S_0 and $\pi^*(S'_1)$. Using that the filter \mathcal{F} is normal, it is not difficult to see that the image of a name σ in HS under an automorphism π^* from \mathcal{G} is also a name in HS, and moreover if some useful tree T witnesses that σ is symmetric, then $\pi(T)$ witnesses that $\pi(\sigma)$ is symmetric. So $\dot{c}_{p_1} \in \text{HS}$ and S_1 witnesses that it is symmetric. Finally, note that $\text{dom}(p_1) \subseteq S_1$. In the next step, we choose some condition in D_0 incompatible with both p_0 and p_1 together with some name in HS it forces to be in $\dot{F}(0)$. We use an automorphism from H_S to transform the witnessing tree into a tree that can be unioned up with S_1 without growing any branches by moving over nodes (outside S) that have new nodes sitting directly above them. We continue this process transfinitely for as long as we can find a condition in D_0 that is incompatible to all previously chosen conditions. At limit stages, we simply take the union of the increasing sequence of trees constructed up to that stage. The process must terminate by some countable ordinal stage α since $\mathbb{P}(\vec{P}^J, <^\omega \omega_1)$ has the ccc. Thus, the final union tree $T_0 = \bigcup_{\xi < \alpha} S_\xi$ must be countable and by construction, since we never allow any branch to grow more than once, T_0 cannot have an infinite branch.

Let's argue that the tree T_0 , because it contains $\bigcup_{\xi < \alpha} \text{dom}(p_\xi)$, witnesses that the mixed name \dot{C}_0 of the names \dot{c}_{p_ξ} for $\xi < \alpha$ over the antichain $\langle p_\xi \mid \xi < \alpha \rangle$ is symmetric. Recall that

$$\dot{C}_0 = \bigcup_{\xi < \alpha} \{(\tau, r) \mid r \leq p_\xi, r \Vdash \tau \in \dot{c}_{p_\xi}, \tau \in \text{dom}(\dot{c}_{p_\xi})\}.$$

Fix an automorphism π point-wise fixing T_0 . It suffices to argue that whenever $(\tau, r) \in \dot{C}_0$, then $(\pi(\tau), \pi(r)) \in \dot{C}_0$. So suppose $(\tau, r) \in \dot{C}_0$ and fix p_ξ witnessing this. Since $r \leq p_\xi$, it follows that $\pi(r) \leq p_\xi$; since $r \Vdash \tau \in \dot{c}_{p_\xi}$, it follows that $\pi(r) \Vdash \pi(\tau) \in \dot{c}_{p_\xi}$; and finally, since \dot{c}_{p_ξ} is symmetric and $\tau \in \text{dom}(\dot{c}_{p_\xi})$, it follows that $\pi(\tau) \in \text{dom}(\dot{c}_{p_\xi})$. Now observe that, since each $\dot{c}_{p_\xi} \in \text{HS}$, we have $\dot{C}_0 \in \text{HS}$.

Next, we let D_1 be the dense set below q of conditions p such that for some name $\dot{c}_p \in \text{HS}$, $p \Vdash \dot{c}_p \in \dot{F}(1)$. We thin out D_1 to a maximal antichain over which we can mix to obtain the desired symmetric name \dot{C}_1 by the same process as above building the new trees as extensions of T_0 , where branches that have extended beyond S are no longer allowed to grow. Once, \dot{C}_1 and T_1 have been obtained, we proceed to obtain \dot{C}_2 and T_2 extending T_1 , and continue in this manner for ω -many steps. The final tree $T = \bigcup_{n < \omega} T_n$ is still countable, cannot have an infinite branch by construction, and witnesses that the canonical name for the sequence of the \dot{C}_n is symmetric. \square

10. A SECOND-ORDER ARITHMETIC MODEL OF $\Sigma_\infty^1\text{-AC} + \neg\Pi_2^1\text{-DC}$

We will now argue that the reals of the symmetric submodel N of the forcing extension $L[G]$ by $\mathbb{P}(\vec{P}^J, <^\omega \omega)$, constructed in Section 9, is a second-order arithmetic model of \mathbb{Z}_2 together with $\Sigma_\infty^1\text{-AC}$ in which $\Pi_2^1\text{-DC}$ fails.

Theorem 10.1. *There is a β -model of second-order arithmetic \mathbb{Z}_2 together with $\Sigma_\infty^1\text{-AC}$ in which $\Pi_2^1\text{-DC}$ fails.*

Proof. Let $G \subseteq \mathbb{P}(\vec{P}^J, <^\omega \omega_1)$ be L -generic and let $N \subseteq L[G]$ be the symmetric submodel of $\mathbb{Z}\mathbb{F} + \text{AC}_\omega + \neg\text{DC}$ constructed in Section 9. Let \mathcal{M} be the model of \mathbb{Z}_2 whose collection of sets is $P(\omega)^N$. Since AC_ω holds in N , it immediately

follows that Σ_∞^1 -AC holds in \mathcal{M} . Thus, it remains to show that Π_2^1 -DC fails in \mathcal{M} . The result will follow if we can show that the tree \mathcal{T} is Π_2^1 -definable in \mathcal{M} . By Theorem 8.2, a sequence $\langle r_0, \dots, r_{n-1} \rangle$ of reals is in the domain of \mathcal{T} if and only if it is L -generic for \mathbb{P}_n . So we need to see that the statement $\langle r_0, \dots, r_{n-1} \rangle$ is L -generic for \mathbb{P}_n is Π_2^1 -expressible in \mathcal{M} . This will suffice because the sequences in \mathcal{T} are ordered simply by extension.

What we would like to say is that all L_α , with α a limit ordinal, should satisfy the following. Given an ordinal β we check whether stage β was nontrivial in the construction of $\vec{P}^J = \langle \mathbb{P}_n^J \mid n < \omega \rangle$. In this case, we check whether $\langle r_0, r_1, \dots, r_{n-1} \rangle$ generates an M_β -generic filter for \mathbb{P}_n^β . Let H_0 consist of all trees in \mathbb{P}_1^β which have the branch r_0 . If H_0 is not M_β -generic, we are done. Otherwise, let H_1 consist of all conditions $p \in \mathbb{P}_2$ such that $p(0) \in H_0$ and r_1 is a branch through $p(1)_{H_0}$. If H_1 is not M_β -generic, we are done. Otherwise, we continue. So $\langle r_0, r_1, \dots, r_{n-1} \rangle$ is \mathbb{P}_n^J -generic if whenever α is a limit ordinal, then L_α satisfies that for every nontrivial stage β in the construction of \mathbb{P}_n^J , $\langle r_0, r_1, \dots, r_{n-1} \rangle$ generates an M_β -generic filter. Now it is not difficult to check that the complexity of this statement is Π_2^1 , because it says that for every set X , if X codes a limit ordinal α , then there is another set Y coding L_α and a set coding the filters H^β needed to verify for every nontrivial stage $\beta \in L_\alpha$ that H^β is M_β -generic. The statement that X codes an ordinal is Π_1^1 and the remaining statements are Σ_1^1 . \square

To produce our β -model of $Z_2 + \Sigma_\infty^1$ -AC in which Π_2^1 -DC fails we used the consistency of (a fragment of) ZFC. We end this section with the following natural open question.

Question 10.2. Does the consistency of $Z_2 + \Sigma_\infty^1$ -AC + $\neg\Pi_2^1$ -DC follow just from the consistency of Z_2 ? Does this implication hold with "consistency" replaced by "existence of a β -model"?

11. REFLECTION CAN FAIL IN MODELS OF ZFC WITHOUT POWERSSET

Let's call the *Reflection Principle* the statement that for every set a and formula $\varphi(\bar{x}, a)$, there is a transitive set model M containing a such that $\varphi(\bar{x}, a)$ is absolute between M and the universe. By the Lévy-Montague reflection, ZFC implies the Reflection Principle, namely every formula is reflected by some V_α . It is natural to wonder whether the Reflection Principle holds in models of ZFC^- , which may not have the V_α -hierarchy. It has been suspected for a long time that the Reflection Principle can fail in such models. The question was first asked by Zarach in [Zar96], and considered again in [GHJ]. We will show that the Reflection Principle fails in the model $H_{\omega_1}^N$, where N is the symmetric submodel of the forcing extension $L[G]$ by $\mathbb{P}(\vec{P}^J, <^\omega \omega_1)$ that we constructed in Section 9.

The argument requires first seeing that the Reflection Principle is equivalent to a version of DC for definable relations. Following [GHJ], let's define that the *Dependent Choice Scheme*, abbreviated DC-scheme, asserts for every formula $\varphi(x, y, z)$ and parameter a that if for every x , there is y such that $\varphi(x, y, a)$ holds, then there is an ω -sequence $\langle x_n \mid n < \omega \rangle$ such that for all n , $\varphi(x_n, x_{n+1}, a)$ holds; in other words, if a definable relation has no terminal nodes, then we can make ω -many dependent choices according to it.

Lemma 11.1 ([GHJ]). *The Reflection Principle is equivalent over ZFC^- to the DC-scheme.*

Proof. First, let's assume that the Reflection Principle holds. Fix a relation $\varphi(x, y, a)$ without terminal nodes. Let M be a transitive set which reflects $\varphi(x, y, a)$ and let R be the set relation on M derived from $\varphi(x, y, a)$. Now fix a well-ordering W of M and use it to define an ω -sequence of choices according to R .

Next, suppose that the DC-scheme holds. Fix a formula $\varphi(\bar{x}, a)$. Observe that given any set A , using collection, we can argue that there is a set $\bar{A} \supseteq A$ which is closed under existential witnesses for subformulas of $\varphi(\bar{x}, a)$ with parameters from A . By taking the transitive closure we can assume that \bar{A} is transitive. So consider the definable relation R which says that A is related to \bar{A} , whenever $A \subseteq \bar{A}$, \bar{A} is transitive, and \bar{A} is closed under existential witnesses for subformulas of $\varphi(\bar{x}, a)$ with parameters from A . We just argued that R has no terminal nodes. Thus, we can make a sequence $\langle M_n \mid n < \omega \rangle$ of dependent choices in R . But then clearly $M = \bigcup_{n < \omega} M_n$ reflects $\varphi(\bar{x}, a)$. \square

Theorem 11.2. *The theory ZFC^- does not imply the Reflection Principle.*

Proof. Let $M = H_{\omega_1}^N$, where N is the ZF-model constructed in Section 9. Clearly $M \models ZF^-$ and since $N \models AC_\omega$, choice holds in M , so $M \models ZFC^-$. Since \mathcal{T} is definable in M (by the argument given in the proof of Theorem 10.1), it follows that the DC-scheme fails in M . Thus, by Lemma 11.1, the Reflection Principle fails as well. \square

REFERENCES

- [Abr84] Uri Abraham. A minimal model for $\neg CH$: iteration of Jensen's reals. *Trans. Amer. Math. Soc.*, 281(2):657–674, 1984.
- [GHJ] Victoria Gitman, Joel David Hamkins, and Thomas A. Johnstone. What is the theory ZFC without power set? To appear in *Mathematical Logic Quarterly*.
- [Jen67] Ronald B. Jensen. Consistency results for models of ZF. *Notices Am. Math. Soc.*, 14:137, 1967.
- [Jen70] Ronald Jensen. Definable sets of minimal degree. In *Mathematical logic and foundations of set theory (Proc. Internat. Colloq., Jerusalem, 1968)*, pages 122–128. North-Holland, Amsterdam, 1970.
- [KL] Vladimir Kanovei and Vassily Lyubetsky. A countable definable set of reals containing no definable elements.
- [Lév70] Azriel Lévy. Definability in axiomatic set theory. II. In *Mathematical Logic and Foundations of Set Theory (Proc. Internat. Colloq., Jerusalem, 1968)*, pages 129–145. North-Holland, Amsterdam, 1970.
- [Sim09] Stephen G. Simpson. *Subsystems of second order arithmetic*. Perspectives in Logic. Cambridge University Press, Cambridge; Association for Symbolic Logic, Poughkeepsie, NY, second edition, 2009.
- [Zar96] Andrzej M. Zarach. Replacement \rightarrow collection. In *Gödel '96 (Brno, 1996)*, volume 6 of *Lecture Notes Logic*, pages 307–322. Springer, Berlin, 1996.

(S.-D. Friedman) KURT GÖDEL RESEARCH CENTER, UNIVERSITY OF VIENNA, WÄHRINGER STRASSE 25 A-1090 VIENNA, AUSTRIA

E-mail address: sdf@logic.univie.ac.at

URL: <http://www.logic.univie.ac.at/~sdf/>

(V. Gitman) THE CITY UNIVERSITY OF NEW YORK, CUNY GRADUATE CENTER, MATHEMATICS PROGRAM, 365 FIFTH AVENUE, NEW YORK, NY 10016

E-mail address: vgitman@nylogic.org

URL: <http://boolesrings.org/victoriagitman>